CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 40, 2011

A

Abe, F., 281
Adamczyk, J. J., 1494
Adler, P. H., 289
Aguirre, C., 1437
Ahn, S. Z., 1239
Alcaide, J., 931
Aliakbarpour, H., 873
Allen, D., 1067
Allen, D. C., 1266
Almeida, R. P. P., 755, 1192, 1595
Aluja, M., 727
Alvarez, J. M., 350
Alyokhin, A., 570
Amézquita, S., 239
Aminatou, B. A., 788
Amore, V., 167
An, C.-J., 1566
April, V., 1123
Arbogast, R. T., 82
Artz, D. R., 1144
Ascher, J. S., 621
Ayres, B. D., 1530
Ayres, M. P., 824, 1007, 1530

B

Bachman, P. M., 1613
Baird, D., 1276
Baldauf, M. W., 639
Balusu, R. R., 1471
Bargues, L., 931
Baril, A., 560
Barraclough, E. I., 1331
Barrera, J. F., 441
Barrette, M., 737
Bayoumy, M. H., 835
Beard, C. E., 289
Becerra, J. X., 1102
Beckman, T. G., 1465
Bel-Kadhi, M. S., 809
Benda, N. D., 1478
Berg, J. V. D., 477
Bergh, J. C., 173, 880
Berkebile, D. R., 184
Berlocher, S. H., 1388
Beuzelin, J. M., 1036
Bextine, B., 23
Bishir, J., 273
Bisseleua, D.h.b., 111
Björkman, C., 333
Black, S. H., 503
Blossey, B., 1074
Blouin, D. C., 1036
Blubaugh, C. K., 1093
Boiteau, G., 94
Boivin, G., 737
Bosch, J., 126
Bottrell, D. G., 42
Boudon-Padieu, E., 1345
Boughton, A. J., 1448
Boutin, C., 560
Bouton, L., 1604
Brandl, D. G., 706
Brehme, C., 204
Brent, C. S., 455, 1554
Bressan, A., 1345
Brévault, T., 484
Brewster, C. C., 173
Brightwell, R. J., 1019
Brodeur, J., 737
Brown, M. W., 42
Brownie, C., 1478
Buergi, L. P., 1370
Burgess, E. P. J., 1331
Burks, C. S., 706
Byeon, Y. W., 431
Byers, J. A., 1210
Bylund, H., 333

C

Caceres, V. A., 1093
Cafaro, M. J., 342
Cai, J. J., 1487
Campos, J. M., 931
Cane, J. H., 614
Carrasco, J. V., 441
Carrière, Y., 484
Carson, D. B., 1613
Caruth, C., 989
Casagrande, R. A., 1427
Cave, R. D., 948
Cha, D. H., 1511
Chantos, J., 324
Charlet, L. D., 15
Chavarría, F., 1111
Chen, M., 496
Chen, S., 505
Chen, Y., 30, 1323, 1405
Cheng, P., 30
Choi, M. Y., 431
Choi, W. I., 1363
Christian, S. J., 1276
Coghlin, P. C., 818
Colpitts, B. G., 94
Cooper, M. L., 1595
Cooper, W. R., 151, 367
Cossé, A., 904
Cottrell, T. E., 1465
Cox, J. E., 1417
Crook, D., 904
Crowder, D. W., 484
Cruz-López, L., 441

D

Daane, K. M., 315, 1595
Daniels, J. C., 1295
Daugherty, M. P., 1192
Dâvalos, A., 1074
Deglow, E. K., 1458
Delphia, C. M., 917
Diaz, R., 1437
Dillwith, J. W., 151
Ditommaso, A., 194
Dosdall, L., 818
Dowd, S. E., 23
Drummond, F. A., 570
Duehl, A. J., 82, 273
Durkee, C. A., 1223

E

Eager, P. T., 1266
Egami, C., 462
Egidi, G., 167
Epstein, D. L., 661
Evans, L. M., 824

F

Fadamiro, H. Y., 1471
Fakhfakh, H., 809
Fantinou, A. A., 597, 1577
Farrell, L. D., 669
Favila, M. E., 239
Feder, J. L., 1353
Fibla, J. M., 931
Fidgen, J. G., 679
Fielding, D. J., 541
Fierke, M. K., 1266
Filella, I., 126
Fisher, R., 204
Flaherty, L., 1200
Fleischer, S. J., 505
Flexner, J. L., 964
Floate, K. D., 818

Fochetti, R., 167
Foley, C. J., 1303
Folgarait, P. J., 342
Frair, J. L., 1266
Frank, D. L., 173
Fraser, I., 904
Frohnapple, K. J., 374
Fu, Q., 749
Fuchs, M., 194
Fusco, R. A., 1417

G

Gaigl, A., 770
Galen, C., 534
Gao, Y., 749
García, F. J. M., 1345
García, M. A. P., 167
García-Marí, F., 931
García-Ruiz, E., 939
Gardner, K. T., 66
Gautam, B. K., 1232
Gautam, S. G., 788
Ge, F., 462
Geden, C. J., 88
Gils, H. A. J. A. V., 770
Giraldeau, L.-A., 737
Gnanvossou, D., 844
Godfrey, L. D., 633, 689
Goergen, G., 844
Gómez, L. E., 770
Goodman, R. M., 1111
Gorsane, F., 809
Goughnour, R. B., 1353
Gould, A., 1131
Gould, F., 1478
Granier, M., 989
Grant, G., 904
Gray, E. W., 1417
Greco, N., 247
Green, G., 120
Grieshop, M. J., 661
Grönberg, L., 333
Gruber, M., 1622
Grubin, S. M., 1410
Grundel, R., 374
Grunwald, S., 893
Guénard, B., 1523
Guenther, D. A., 66
Guo, Y., 1323
Gut, L. J., 661
Gutierrez, A. P., 755
Guy, M., 560

H

Haack, R. A., 1363
Hahn, D. A., 1295
Hail, D., 23
Hain, F. P., 273
Halima, A. B., 809
Hall, C. L., 669
Hamilton, G. C., 231, 1131
Hance, T., 737
Hanks, L. M., 1388
Hannoufa, A., 1622
Harmon, J. P., 103
Harris, M. K., 889
Hartfield, E. A., 889
Hartke, T. R., 1554
Hatle, J. D., 157
He, H., 1405
He, L., 1487
Hejazi, M. J., 1253
Helali, H., 1253
Hellmich, R. L., 496
Henderson, G., 1232
Henne, D. C., 73
Herms, D. A., 648
Hesler, S. P., 1511
Heuberger, S., 484
Higbee, B. S., 706
Hoddle, M., 755
Hoeting, J. A., 654
Hofstetter, R. W., 824
Hogsette, J. A., 88
Holdcraft, R., 412
Hooks, C. R. R., 391
Hortoa, D. L., 1465
Hotchkiss, M. W., 59
Howard, D. R., 669
Hsu, C. L., 1144
Hu, L., 743
Hu, Yi, 516
Hu, Y., 1
Hudson, I., 589

I

Iburg, J. P., 1417
Ikemoto, T., 462
Iranipour, S., 1253
Irvin, N. A., 755
Isaacs, R., 399, 697, 1177
Ives, A. R., 103

J

Jablonski, S. E., 1604
James, L.-T., 1554
James, R. R., 782
Janzen, D. H., 1111
Jean, R. P., 374
Jennings, E. M., 669
Jensen, L. B. M., 1458
Jia, B., 1487
Jiang, C.-J., 1566, 1613
Jiang, X., 516
Jo, Y.-K., 589
Joern, A., 979
Johnson, M. W., 315
Johnson, S. J., 73

K

Kampouris, S., 1541
Kanzaki, N., 281
Kaplan, I., 1093
Karamaouna, F., 1541
Karimzadeh, R., 1253
Kaufman, P. E., 88
Kay, F. M., 605
Kean, A. M., 1331
Kemp, W. P., 917
Khalifa, M. B., 809
Kim, D., 880
Kim, J. H., 431
Klepzig, K. D., 824
Klingeman, W. E., 1397
Kluger, E. C., 1388
Knight, A. L., 420
Kok, L. T., 324
Kollmann, J., 1183
Kolokytha, P. D., 597, 1577
Kontodimas, D. C., 1541
Koppenhöfer, A. M., 412
Kreutzian, M., 1051
Kroschel, J., 217
Kruger, M., 477
Krupke, C. H., 1585

L

Lai, F., 749
Laing, M. D., 52
Lamb, A. B., 324
Lambert, B. D., 1572
Langor, D. W., 797
Lapointe, S. L., 1437
Larson, J., 1093
Larsson, L., 333
Lashomb, J., 1131
Latty, T. M., 864
Lauzière, I., 23
Laws, A. N., 979
Lee, C.-Y., 470
Lee, J.-S., 1363
Lei, C., 1
Lei, Z., 132
Lemay, M. A., 904
Leskey, T. C., 94, 173
Lester, P. J., 1276
Levine, S. L., 1613
Li, D.-C., 140

Li, Xiang, 1622
Li, Xue, 132
Li, Yongjian, 30
Li, Yunhe, 496, 749, 1323
Liburd, O. E., 893
Light, D. M., 420
Lightfoot, D. C., 1051
Linn, C. E. J., 1511
Littlefield, K. A., 1572
Liu, T.-X., 140
Liu, X., 496
Liu, Zhongfang, 449
Liu, Zhuorong, 749
Llusià, J., 126
Loeb, G. M., 1511
Lombardero, M. J., 1007
López-Guillén, G., 441
Lowery, D. T., 1458
Lu, W., 1487
Lucky, A., 1523
Luo, J., 749
Luo, L., 516
Lyons, D. B., 904

M

Mace, W. J., 639
Mafra-Neto, A., 1494
Magee, D., 714, 904
Magnuson, T. S., 669
Malik, N., 1494
Mallis, R. E., 1378
Malo, E. A., 441
Malone, L. A., 1331
Manandhar, R., 391
Mangan, R., 1494
Manosalva, L., 399
Marchal, C., 989
Marco, V., 939
Marcum, D. B., 689
Marfetán, J. A., 342
Markussen, B., 1183
Martin, P. A., 560
Martínez, Y., 989
Martínez-Ferrer, M. T., 931
Marzal, C., 931
Masuya, H., 281
Mathews, C. R., 42
Matsuda, T., 204
Mayo, P. D., 714, 904
McAvoy, T. J., 324
McCabe, S. K., 560
McCaffrey, J., 534
McCreadie, J. W., 289
McCullough, D. G., 357, 1239
McGraw, B. A., 412
Medina, R. F., 889
Meer, R. K. V., 1276
Meikle, W. G., 1494
Meinke, L. J., 964
Meisner, M. H., 103
Meloche, F., 94
Mendoza, M. G., 549
Merrill, S. C., 654
Mészáros, A., 1036
Meyer, J. R., 880
Miao, J., 743
Middleton, M. C., 1595
Miller, J. R., 661
Miller, R., 52
Miller, S. A., 614
Mills, N. J., 1370
Miranda, F., 333
Mitrovich, M., 204
Mohammadi, S. A., 1253
Moharramipour, S., 1546
Montemayor, C. O., 948
Moore, D. I., 1051
Moore, L. A. S., 260
Mora, M., 1111
Morales-Ramos, J. A., 1285
Morton, P. K., 1303
Mostafa, A. M., 1458
Muilenburg, V. L., 648
Muir, J. P., 1572
Mujica, N., 217
Muñiz, Y., 989
Murphy, A. F., 1585
Mutis, A., 399
Mwamburi, L. A., 52

N

Nault, B. A., 194, 1144
Navarro-Campos, C., 931
Negro, M., 1081
Nielsen, A. L., 231
Nielsen, D. G., 648
Noblet, R., 1417
Noge, K., 1102
Nowatzki, T. M., 964
Nyrop, J. P., 381

O

Ode, P. J., 15
Oetting, R. D., 132
Okabe, K., 281
Oleiro, M., 605
Olivier-Espejel, S., 1102
Olivo-Martínez, A., 549
O'Neal, M. E., 260
O'Neill, K. M., 917
O'Neill, R. P., 917
Onstad, D. W., 964
Opit, G. P., 788
Opp, S. B., 315
Ortega, F., 399
Ostrem, J., 496
Othman, A. S., 470
Overholt, W. A., 1437

P

Paine, T. D., 36
Palestrini, C., 1081
Pan, Z., 964
Papadoulis, G. T., 597, 1577
Paradise, M. S., 1613
Paranhos, B., 727
Pardee, G. L., 581
Pardo, F., 399
Parmenter, R. R., 1051
Parra, L., 399
Parry, D., 955
Patt, J. M., 1494
Pavis, C., 989
Pavlovic, N. B., 374
Peairs, F. B., 654
Pedersen, A. B., 633
Pemberton, R. W., 1448
Peng, Y., 496, 749, 1323
Peñuelas, J., 126
Pérez-Moreno, I., 939
Perfecto, I., 999
Perich, F., 399
Peterschmitt, M., 989
Philpott, S. M., 581
Pierce, S. J., 1239
Pinto-Tomás, A. A., 1111
Pinzon, J., 797
Pitt, D., 904
Pitzer, J. B., 88
Poland, T. M., 357, 1239
Ponti, L., 755
Post, K. H., 955
Pureswaran, D., 1200
Puterka, G. J., 151

Q

Qin, A. Z., 1487
Quiring, D. T., 1200
Quiroz, A., 399
Qureshi, J. A., 1503

R

Rashed, A., 1192
Rawi, C. S. M., 873
Reagan, T. E., 1036
Reeves, J. L., 855
Reilly, C. C., 59
Reisig, D. D., 689
Rendón, P. A., 315

Rensburg, J. B. J. V., 477
Rhodes, E. M., 893
Richmond, D. S., 639, 1093
Rieske, L. K., 1378
Robertson, M. P., 1123
Rocca, M., 247
Rochester, C., 204
Rodriguez-Saona, C., 412
Roelofs, W. L., 1511
Rojas, J. C., 441
Rojas, M. G., 1285
Rolando, A., 1081
Roltsch, W. J., 1370
Romeis, J., 496
Rosengaus, R. B., 1554
Ross, D. W., 864, 1410
Rosskopf, E., 1437
Ryall, K. L., 679, 904

S
Sabree, Z. L., 1102
Sadof, C. S., 1093
Salinas-Moreno, Y., 549
Salom, S. M., 324
Sampson, B. J., 614
Sanders, N. J., 1397
Sanderson, J. P., 381
Sappington, T. W., 516
Saunders, M. C., 505
Scarr, T., 904
Schal, C., 1478
Schemerhorn, B. J., 1303
Schmidt, N. P., 260
Seco, R., 126
Seiler, G. J., 15
Sétamou, M., 1494
Sewell, G., 570
Sforza, R., 1595
Shakya, K., 788
Shapiro-Ilan, D. I., 59, 1285
Shearer, P. W., 231
Shelton, A. M., 194, 496
Shi, P., 462
Shipp, L., 381
Sholes, O. D. V., 777
Showler, A. T., 1036
Shufran, K. A., 1317
Shumate, A. M., 1530
Siegel, J. P., 706
Siegert, N. W., 1239
Sigsgaard, L., 1183
Silk, P. J., 714, 904
Silverman, J., 1019
Simelane, D. O., 1123
Simon, H. M., 1111
Sittenfeld, A., 1111
Sivinski, J., 727
Skou, A.-M. T., 1183
Smith, E. A., 194
Smith, R. J., 669
Smitley, D. R., 589
Snyder, W. E., 8
Sopow, S., 714
Soroka, J., 1622
Soudi, S., 1546
Spence, J. R., 797
Spurgeon, D. W., 367, 455
Srinivasan, R., 350
Stanley, B. H., 964
Stansly, P. A., 1503
Stathas, G. J., 1541
Stauffer, T. W., 157
Stelinski, L. L., 661
Storch, R. H., 570
Straw, N. A., 120
Stringer, L. D., 1276
Stuhl, C., 727
Stump, A. D., 1604
Suckling, D. M., 1276
Sun, Y., 462
Sweeney, J. D., 714, 904, 1200
Szendrei, Z., 412

T
Tabashnik, B. E., 484
Takada, M., 1027
Takagi, M., 431
Taki, H., 281
Takizawa, T., 8
Talley, J., 788
Tan, J., 1613
Tanaka, R., 281
Taylor, D. B., 184
Teal, P. E., 82, 727, 1511
Teale, S. A., 1530
Tedders, W. L., 1285
Thompson, D. C., 66
Tian, M. Y., 1487
Tillman, P. G., 303, 1159
Timperio, A. M., 167
Tindo, M., 844
Tluczek, A. R., 357
Tobin, P. C., 505
Toennisson, T. A., 1397
Tomberlin, J. K., 30, 1572
Tooker, J. F., 1388
Trimble, R. M., 420
Tuda, M., 431
Turgeon, J. J., 679
Turschak, G., 204

U
Ubero-Pascal, N., 167
Uffman, J. P., 1613
Ugine, T. A., 381
Uma, D. B., 1223
Umaharan, P., 989
Uribe-Lorío, L., 1111

V
Vail, K. M., 1397
Vandermeer, J. H., 999, 1067
Vayssières, J.-F., 844
Vidal, S., 111
Vieira, L. C., 324
Vincent, C., 94

W
Wadsworth, N. K., 669
Wallin, K. F., 864, 1410
Walsh, D. B., 1595
Walter, C., 1331
Walter, S. M., 654
Walton, N. J., 697, 1177
Walton, V. M., 1595
Wan, F.-H., 140
Wang, H., 132
Wang, J.-J., 140
Wang, K., 381
Wang, Q., 1487
Wang, X., 1
Wang, X.-G., 315
Warren, V., 1295
Washitani, I., 1027
Watson, J. C., 621
Way, M. O., 1036
Weed, A. S., 1427
Wei, C., 1405
Wei, S., 1622
Weiss, M. R., 1223
Wheeler, G. S., 605, 1437
Whitman, D. W., 157
Wicker, E., 989
Wilder, J. A., 1604
Williams, D. T., 120
Wilson, L. T., 1036
Wolf, A. T., 621
Wooley, S. C., 36
Wraight, S. P., 381
Wu, G.-M., 737
Wu, J., 714
Wu, K., 449, 1323
Wu, Y., 743
Wyatt, R. D., 1417
Wyckhuys, K. A. G., 449

X

Xu, J., 1487
Xu, N., 1622
Xu, Q., 743
Xu, W., 743

Y

Yang, C., 1494
Yang, Z., 30
Yeap, B.-K., 470
Yede,, 111
Yee, W. L., 1353
Yitbarek, S., 1067
Yokoyama, V. Y., 315
Yoshioka, A., 1027
Yu, G., 30
Yu, Z., 743

Z

Zaviezo, T., 1595
Zebarjadi, A., 1622
Zhang, A., 880, 1511
Zhang, B.-C., 1566
Zhang, G.-F., 140
Zhang, J., 1131
Zhang, L., 516
Zhang, M., 541
Zhang, Q.-W., 1566
Zhang, Y., 1323, 1405
Zhao, Z.-W., 1566
Zhu, F., 1
Zúñiga, G., 549

CUMULATIVE KEY WORD INDEX
Environmental Entomology
Volume 40, 2011

A

Área de Conservación Guanacaste,
  insect-microbe interactions, 1111
  midgut microbiota, 1111
    Rothschildia lebeau, 1111
    tropical caterpillar, 1111
Abagrotis orbis,
  climbing cutworms, 1458
  feeding preferences, 1458
  grapevine, 1458
  host plant, 1458
Acrididae,
  bottom-up forces, 541
  Orthoptera, 541
  resource limitation, 541
  subarctic, 541
active hunters,
  habitat structure, 1378
  predators, 1378
  Tsuga canadensis, 1378
  web builders, 1378
additive diversity partitioning,
  Generalized Linear Models, 1081
  IndVal, 1081
  overgrazing, 1081
  Scarabaeidae, 1081
Adelges tsugae, 324, 864, 1410
  biological control, 324, 864, 1410
  hemlock woolly adelgid, 324
  host finding behavior, 864
  Host range, 324
  Laricobius osakensis, 324
  Leucopis argenticollis, 1410
  Leucopis piniperda, 1410
  predator-prey interactions, 864
  prey suitability, 1410
  Tsuga spp., 864
adult distribution,
  codling moth, 661
  diel behavior, 661
  mating disruption, 661
  vacuum, 661
aggregation,
  autocorrelation, 689
  pest management, 689
  Semivariogram, 689
  Taylor's power law, 689
Agrilus anxius,
  biogeography, 648
  coevolution, 648
  host plant resistance, 648
  wood-borers, 648
Agrilus planipennis, 904, 1239
  double-decker traps, 1239
  Fraxinus, 1239
  girdled trees, 1239
  green trap, 904
  kairomone, 904
  lactone, 904
  Manuka oil, 1239
  pheromone, 904
agro-ecosystem,
  Andrenidae, 621
  apparent competition, 1027
  apple orchards, 621
  Italian ryegrass, 1027
  landscape ecology, 621
  moth diversity, 560
  native bees, 621
  organic farming, 560
  plant diversity, 560
  plant introduction, 1027
  woody hedgerow, 560
agroforestry,
  Azteca instabilis, 581
  nonconsumptive effects, 581
  Pseudacteon, 581
  trait-meditated interactions, 581
alatae,
  host utilization, 350
  intraguild competition, 350
  Macrosiphum euphorbiae, 350
  Myzus persicae, 350
alfalfa,
  alfalfa leafcutting bees, 782
  bees, 782
  chalkbrood, 782
  disease transmission, 782
alfalfa leafcutting bee,
  alfalfa, 782
  bees, 782
  chalkbrood, 782
  disease transmission, 782
  lipids, 917
  Megachile rotundata, 917
  pollinators, 917
  thermal biology, 917
alien species,
  Bemisia tabaci, 140
  co-occurrence, 140
  interspecific interaction, 140
  Trialeurodes vaporariorum, 140
alkaloids,
  ergovaline, 639
  N-acetyl norloline, 639
  Neotyphodium, 639
  peramine, 639
Allium cepa,
  population ecology, 194
  Thrips tabaci, 194
  Tospovirus, 194
almonds,
  Amyelois transitella, 706
  egg trap, 706
  monitoring, 706
  Prunis dulcis, 706
alternate hosts, 633, 1036
  Diabrotica undecimpunctata undecimpunctata, 633
  Diatraea saccharalis (F.), 1036
  Eoreuma loftini (Dyar), 1036
  larval development, 633
  Mexican rice borer, 1036
  sugarcane borer, 1036
alternative host,
  gene-for-gene interaction, 1303
  landscape genetics, 1303
  microsatellites, 1303
  wheat, 1303
altitudinal range,
  behavioral thermoregulation, 534
  colonization, 534
  Formica neorufibarbis, 534

nest site preference, 534
Amazon,
Atta sexdens, 770
Colombia, 770
demography, 770
palatable forage hypothesis, 770
Amyelois transitella,
almonds, 706
egg trap, 706
monitoring, 706
Prunis dulcis, 706
Anastrepha,
Bactrocera, 727
Ceratitis, 727
Diachasmimorpha longicaudata, 727
para-ethylacetophenone, 727
Andrenidae,
agro-ecosystem, 621
apple orchards, 621
landscape ecology, 621
native bees, 621
annual bluegrass weevil,
EAG, 412
GC-MS, 412
green leaf volatiles, 412
Listronotus maculicollis, 412
ant community,
coastal sage scrub, 204
southern California, 204
species diversity, 204
wildfire, 204
ant mosaic,
ants, 999
pastures, 999
scramble competition, 999
self-organization, 1067
spatial pattern, 999, 1067
ant pests,
biological control, 342
fungi-fungi interactions, 342
induction of conidiation, 342
leaf-cutter ants, 342
antimicrobial peptides,
competition, 669
Nicrophorus, 669
parental care, 669
secretions, 669
ants, 42, 999, 1019
ant mosaic, 999
extrafloral nectaries, 42
habitat modification, 1019
invasive species, 1019
microclimate, 1019
niche expansion, 1019
oriental fruit moth, 42
pastures, 999
peach, 42
plant defense, 42
scramble competition, 999
spatial pattern, 999
aphids, 570, 1541
coexistence, 570
path analysis, 570
population regulation, 570
predator, 1541
pumpkin, 1541
rearing, 1541
thermal thresholds, 1541
Aphis gossypii,
barrier crop, 391
Myzus persicae, 391
nonpersistent virus, 391
virus sink, 391
Aphrophoridae,
Cicadellidae, 1131
Clastopteridae, 1131
Membracidae, 1131
Xylella fastidiosa, 1131
apparent competition,
agroecosystem, 1027
Italian ryegrass, 1027
plant introduction, 1027
apple,
geostatistics, 173
median polishing, 173
residuals, 173
Synanthedon scitula, 173
apple orchards,
agro-ecosystem, 621
Andrenidae, 621
landscape ecology, 621
native bees, 621
Arabidopsis,
beta-ionone, 1622
carotenoid cleavage dioxygenase one over-expression, 1622
resistance to crucifer flea beetles, 1622
Arachnida,
Coleoptera, 1523
Formicidae, 1523
Micro-Hymenoptera, 1523
Winkler extractor, 1523
Arrhenius plot,
enzyme inactivation, 462
intrinsic optimum temperature, 462
Sharpe-Schoolfield model, 462
tangent equation, 462
Asian citrus psyllid,
biological control, 1503
Candidatus liberibacter asiaticus, 1503
Huanglongbing, 1503
life table, 1503
Atta sexdens,
Amazon, 770
Colombia, 770
demography, 770
palatable forage hypothesis, 770
attack density,
dieback, 1363
Korean oak wilt, 1363
Mongolian oak, 1363
Platypus koryoensis, 1363
attack rate,
functional response, 835
handling time, 835
Homalotylus flaminius, 835
numerical response, 835
autocorrelation,
aggregation, 689
pest management, 689
Semivariogram, 689
Taylor's power law, 689
Azteca instabilis,
agroforestry, 581
nonconsumptive effects, 581
Pseudacteon, 581
trait-meditated interactions, 581

B

β-farnesene,
farnesol, 420
Monitoring, 420
pear ester, 420
(Z)-3-hexenyl acetate, 420
β-ionone,
Arabidopsis, 1622
carotenoid cleavage

dioxygenase one over-expression, 1622
resistance to crucifer flea beetles, 1622
Bacillus subtilis,
chicken manure, 30
Hermetia illucens, 30
waste management, 30
Bacillus thuringiensis,
feeding behavior, 484
refuge strategy, 484
simulation modeling, 484
transgenic crops, 484
Bacillus thuringiensis subsp. *israelensis*,
biological control, 52
diatoms, 1417
insecticidal crystalline proteins, 1417
Musca domestica, 52
poultry manure, 52
seston, 1417
Simulium vittatum, 1417
Bactrocera dorsalis complex,
invasive fruit fly, 844
polyphagous, 844
quarantine pest, 844
Tephritidae, 844
Bactrocera oleae (Rossi),
Bactrocera oleae (Rossi), 315
Larval parasitoid, 315
Olea europaea L., 315
Bactrocera,
Anastrepha, 727
Ceratitis, 727
Diachasmimorpha longicaudata, 727
para-ethylacetophenone, 727
bark beetles,
Dendroctonus rhizophagus, 549
geographical distribution, 549
predictive models, 549
bark moisture content,
colonization, 1487
Glenea cantor, 1487
host selection, 1487
oviposition attractant, 1487
barrier crop,
Aphis gossypii, 391
Myzus persicae, 391
nonpersistent virus, 391
virus sink, 391
Batesian mimicry,
imperfect mimicry, 1223
myrmecomorphy, 1223
Peckhamia, 1223
Salticidae, 1223
Beauveria bassiana,
fungicide, 59
genetic improvement, 59
Metarhizium brunneum, 59
resistance, 59
bees, 374, 782
alfalfa, 782
alfalfa leafcutting bees, 782
bowl trapping, 374
chalkbrood, 782
disease transmission, 782
inventory, 374
netting, 374
sampling, 374
behavior, 880, 1285
dietary self-selection, 1285
insect rearing, 1285
nutrition, 1285
plant-insect interactions, 880
roots, 880
Vitacea polistiformis, 880
yellow mealworm, 1285
behavioral thermoregulation,
altitudinal range, 534
colonization, 534
Formica neorufibarbis, 534
nest site preference, 534
Bemisia argentifolii,
Eretmocerus eremicus, 36
Glomus etunicatum, 36
multi-trophic interactions, 36
silverleaf whitefly, 36
Bemisia tabaci, 132, 140, 809
alien species, 140
co-occurrence, 140
cold hardiness, 132
HSP genes, 132
interspecific interaction, 140
rapid cold hardening, 132
secondary symbionts, 809
Trialeurodes vaporariorum, 140
Tunisia, 809
binomial sampling,
classification sampling, 381
flower taps, 381
operating characteristic, 381
visual inspection, 381
bioassay,
chemical attraction, 82
disturbance, 82
Tribolium castaneum, 82
biocontrol,
colonization, 1074
Galerucella, 1074
Lythrum salicaria, 1074
matrix habitat, 1074
biodiversity,
biological control, 8
ecosystem functioning, 8
lady beetle, 8
predatory bugs, 8
biogeography,
Agrilus anxius, 648
coevolution, 648
host plant resistance, 648
wood-borers, 648
biological control, 8, 52, 103, 217, 260, 324, 342, 864, 1093, 1123, 1410, 1448, 1503
Adelges tsugae, 324, 864, 1410
ant pests, 342
Asian citrus psyllid, 1503
Bacillus thuringiensis variety *israelensis*, 52
biodiversity, 8
Candidatus liberibacter asiaticus, 1503
cannibalism, 103
Carabidae, 1093
Coccinellidae, 103
competitive interaction, 1123
Cyclotrachelus sodalis, 1093
ecosystem functioning, 8
ecosystem service, 260
fungi-fungi interactions, 342
hemlock woolly adelgid, 324
host finding behavior, 864
host quality, 1123
Host range, 324
Huanglongbing, 1503
induction of conidiation, 342
intraguild predation, 103
IPM, 1093
lady beetle, 8
Laricobius osakensis, 324
larval tracks, 103
leaf-cutter ants, 342
leaf-galls, 1448
Leucopis argenticollis, 1410

Leucopis piniperda, 1410
life table, 1503
Liriomyza huidobrensis, 217
Liriomyza sativae, 217
Lygodium microphyllum, 1448
multiple release, 1123
Musca domestica, 52
omnivory, 1093
parasitoids, 217
population dynamics, 1123
poultry manure, 52
predator-prey interactions, 864
predatory bugs, 8
prey suitability, 1410
tritrophic interaction, 260
Tsuga spp., 864
biomass,
carrion removal rates, 239
forest fragmentation, 239
necrophagous beetles, 239
Scarabaeinae, 239
biosafety,
insect host-parasitoid, 1331
leafy gene, 1331
*np-ENDITAL*II selectable marker gene, 1331
risk assessment, 1331
biotypes,
cytochrome c, 1317
elongation factor 1-α, 1317
greenbug, 1317
host-plant resistance, 1317
blueberry,
conservation biocontrol, 697
fruit, 697
habitat management, 697
integrated pest management (IPM), 697
body size, 1, 931
development time, 1
genotype-temperature interactions, 1
host, 931
Mediterranean fruit fly, 931
sterile insect technique, 931
temperature, 1, 931
body size ratio,
foraging, 737
handling time, 737
temperature-size rule, 737
Bombus impatiens,
Cucurbita spp., 1144
Peponapis pruinosa, 1144
pollination, 1144
bottom-up forces,
Acrididae, 541
Orthoptera, 541
resource limitation, 541
subarctic, 541
bowl trapping,
bees, 374
inventory, 374
netting, 374
sampling, 374
Brazilian pepper,
exocrine gland, 605
host specificity testing, 605
weed biological control, 605
brown stink bug,
cotton, 1159
peanut, 1159
Southern green stink bug, 1159
stink bug distribution, 1159
Bt corn,
female longevity, 1585
head capsule width, 1585
male investment, 1585
stable isotopes, 1585
Bt protein expression,
control efficacy, 1323
ELISA, 1323
transgenic rice, 1323
Buprestidae,
Fraxinus, 679
invasive pest, 679
sample unit, 679
sampling technique, 679
Burkholderia,
Coreidae, 1102
crypts, 1102
gut symbiont, 1102
Wolbachia, 1102

C

Camponotus japonicus,
16S rRNA-restriction fragment-length polymorphism, 1405
gut bacteria, 1405
Candidatus liberibacter asiaticus,
Asian citrus psyllid, 1503
biological control, 1503
Huanglongbing, 1503
life table, 1503
cannibalism,
biological control, 103
Coccinellidae, 103
intraguild predation, 103
larval tracks, 103
canola pest,
MLST, 818
multilocus sequence typing, 818
symbiont, 818
Wolbachia, 818
Carabidae,
biological control, 1093
Cyclotrachelus sodalis, 1093
IPM, 1093
omnivory, 1093
carotenoid cleavage dioxygenase one over-expression,
β-ionone, 1622
Arabidopsis, 1622
resistance to crucifer flea beetles, 1622
carrier specificity,
distribution, 281
Lucanidae, 281
nematode, 281
phoresy, 281
carrion removal rates,
biomass, 239
forest fragmentation, 239
necrophagous beetles, 239
Scarabaeinae, 239
catechin,
cows, 1572
goats, 1572
Musca domestica, 1572
cellular automata,
county pattern, 273
forest damage, 273
southern pine beetle, 273
Cerambycidae, 714, 1200
cinnamopterum, 714
contact pheromone, 714
development time, 1200
fuscum, 714
Tetropium, 714
tree condition, 1200

tree moisture stress, 1200
Ceratitis,
Anastrepha, 727
Bactrocera, 727
Diachasmimorpha longicaudata, 727
para-ethylacetophenone, 727
chalkbrood,
alfalfa, 782
alfalfa leafcutting bees, 782
bees, 782
disease transmission, 782
chemical attraction,
bioassay, 82
disturbance, 82
Tribolium castaneum, 82
chicken manure,
Bacillus subtilis, 30
Hermetia illucens, 30
waste management, 30
Chrysomelidae,
host specificity, 1427
nontarget effects, 1427
prerelease impact, 1427
swallow-worts, 1427
Chrysoperla sinica,
migration, 449
ovarian development, 449
tethered-flight, 449
Cicadellidae,
Aphrophoridae, 1131
Clastopteridae, 1131
Membracidae, 1131
Xylella fastidiosa, 1131
cinnamopterum,
Cerambycidae, 714
contact pheromone, 714
fuscum, 714
Tetropium, 714
classification sampling,
binomial sampling, 381
flower taps, 381
operating characteristic, 381
visual inspection, 381
Clastopteridae,
Aphrophoridae, 1131
Cicadellidae, 1131
Membracidae, 1131
Xylella fastidiosa, 1131
climate change, 505, 777, 824
climate variation, 777
community structure, 824
extreme temperatures, 777
grape berry moth, 505
greenhouse gas emissions, 505
indirect interactions, 824
phenology, 505
physiological ecology, 824
Pinus edulis, 777
species interactions, 824
voltinism, 505
climate variation,
climate change, 777
extreme temperatures, 777
Pinus edulis, 777
climbing cutworms,
Abagrotis orbis, 1458
feeding preferences, 1458
grapevine, 1458
host plant, 1458
co-occurrence,
alien species, 140
Bemisia tabaci, 140
interspecific interaction, 140
Trialeurodes vaporariorum, 140
coastal sage scrub,
ant community, 204
southern California, 204
species diversity, 204
wildfire, 204
Coccinellidae,
Biological control, 103
cannibalism, 103
intraguild predation, 103
larval tracks, 103
cocoon,
frass, 126
Monodontomerus aeneus, 126
Osmia cornuta, 126
volatiles, 126
codling moth,
adult distribution, 661
diel behavior, 661
mating disruption, 661
vacuum, 661
coevolution,
Agrilus anxius, 648
biogeography, 648
host plant resistance, 648
wood-borers, 648
coexistence,
aphids, 570
path analysis, 570
population regulation, 570
coffee berry borer,
degeneration, 441
flight muscles, 441
Hypothenemus hampei, 441
cold hardiness, 132, 1546
Bemisia tabaci, 132
diapause, 1546
elm leaf beetle, 1546
HSP genes, 132
rapid cold hardening, 132
supercooling point, 1546
Xanthogaleruca luteola, 1546
Coleoptera,
Arachnida, 1523
Formicidae, 1523
Micro-Hymenoptera, 1523
Winkler extractor, 1523
Colombia,
Amazon, 770
Atta sexdens, 770
demography, 770
palatable forage hypothesis, 770
colonization, 534, 1074, 1487
altitudinal range, 534
bark moisture content, 1487
behavioral thermoregulation, 534
biocontrol, 1074
Formica neorufibarbis, 534
Galerucella, 1074
Glenea cantor, 1487
host selection, 1487
Lythrum salicaria, 1074
matrix habitat, 1074
nest site preference, 534
oviposition attractant, 1487
color response,
Elatobium abietinum, 120
flight activity, 120
Sitka spruce, 120
sticky traps, 120
community ecology,
habitat structure, 1397
landscaping, 1397
odorous house ant, 1397
urban, 1397
community structure,
climate change, 824
indirect interactions, 824
physiological ecology, 824
species interactions, 824

competition,
antimicrobial peptides, 669
Nicrophorus, 669
parental care, 669
secretions, 669
competitive interaction,
biological control, 1123
host quality, 1123
multiple release, 1123
population dynamics, 1123
computer simulation in three dimensions,
mass trapping, 1210
mating disruption, 1210
mean flight height, 1210
movement algorithms, 1210
conifer pest,
invasive insect, 1266
native parasitoids, 1266
pine mortality, 1266
conservation biocontrol, 697, 1177
blueberry, 697
flowers, 1177
fruit, 697
habitat management, 697, 1177
integrated pest management (IPM), 697
native plants, 1177
natural enemies, 1177
consumption rate,
generalist insect predators, 333
killing rate, 333
natural enemies, 333
spiders, 333
contact pheromone,
Cerambycidae, 714
cinnamopterum, 714
fuscum, 714
Tetropium, 714
control efficacy,
Bt protein expression, 1323
ELISA, 1323
transgenic rice, 1323
copper binding proteins,
insects, 167
nano-RP-HPLC–ESI–MS, 167
Coptotermes formosanus,
moisture, 1232
mortality, 1232
starvation, 1232
Coptotermes gestroi,
gene flow, 470
microsatellite markers, 470
population structure, 470
Coreidae,
Burkholderia, 1102
crypts, 1102
gut symbiont, 1102
Wolbachia, 1102
cotton, 367, 1159
brown stink bug, 1159
feeding behavior, 367
Gossypium, 367
Lygus, 367
peanut, 1159
Southern green stink bug, 1159
stink bug distribution, 1159
western tarnished plant bug, 367
cotton aphid,
fecundity, 431
lower developmental threshold, 431
temperature requirement, 431
thermal constant, 431
county pattern,
cellular automata, 273
forest damage, 273
southern pine beetle, 273
cows,
catechin, 1572
goats, 1572
Musca domestica, 1572
crypts,
Burkholderia, 1102
Coreidae, 1102
gut symbiont, 1102
Wolbachia, 1102
Cuba,
Dominican Republic, 989
Guadeloupe, 989
Martinique, 989
Trinidad, 989
Cucurbita spp.,
Bombus impatiens, 1144
Peponapis pruinosa, 1144
pollination, 1144
Cucurbitaceae,
fruit set, 614
fruit size, 614
oligolecty, 614
pollination efficiency, 614
Curculionidae,
insect conservation, 1388
phytophagy, 1388
plant-insect interaction, 1388
Cyclotrachelus sodalis,
biological control, 1093
Carabidae, 1093
IPM, 1093
omnivory, 1093
Cylindrocopturus adspersus,
native sunflowers, 15
native versus cultivated, 15
parasitism, 15
cytochrome c,
biotypes, 1317
elongation factor 1-α, 1317
greenbug, 1317
host-plant resistance, 1317
cytochrome P450,
detoxification, 1604
Drosophila, 1604
mycophagy, 1604

D

decapitating fly,
diffusion model, 73
dispersal, 73
Solenopsis invicta, 73
defenses,
natural enemies, 1007
nutritional quality, 1007
outbreak, 1007
Scolytinae, 1007
degeneration,
coffee berry borer, 441
flight muscles, 441
Hypothenemus hampei, 441
degree-days,
Epiphyas postvittana, 1370
invasive species, 1370
phenology, 1370
demography,
Amazon, 770
Atta sexdens, 770
Colombia, 770
palatable forage hypothesis, 770
Dendroctonus rhizophagus,
bark beetles, 549
geographical distribution, 549
predictive models, 549

detection,
food preference, 1276
foraging distance, 1276
model, 1276
surveillance, 1276
detoxification,
cytochrome P450, 1604
Drosophila, 1604
mycophagy, 1604
development, 597, 1577
food, 1577
life table, 597
Phytoseiidae, 597
pollen., 597
temperature, 1577
Typhlodromus athenas, 597, 1577
development time, 1, 1200
body size, 1
Cerambycidae, 1200
genotype-temperature interactions, 1
temperature, 1
tree condition, 1200
tree moisture stress, 1200
developmental rate,
field validation, 939
thermal threshold, 939
variable temperature, 939
Xylotrechus arvicola, 939
developmental thresholds,
population growth, 788
stored grain, 788
stored products, 788
developmental time,
predation, 948
spined soldier bug, 948
yellow margined leaf beetle, 948
Diabrotica undecimpunctata undecimpunctata,
alternate hosts, 633
larval development, 633
Diachasmimorpha longicaudata,
Anastrepha, 727
Bactrocera, 727
Ceratitis, 727
para-ethylacetophenone, 727
diapause, 455, 1546
cold hardiness, 1546
elm leaf beetle, 1546
Lygus hesperus, 455
rearing, 455
supercooling point, 1546
western tarnished plant bug, 455
Xanthogaleruca luteola, 1546
diatoms,
Bacillus thuringiensis subsp. *israelensis*, 1417
insecticidal crystalline proteins, 1417
seston, 1417
Simulium vittatum, 1417
Diatraea saccharalis (F.),
alternate hosts, 1036
Eoreuma loftini (Dyar), 1036
Mexican rice borer, 1036
sugarcane borer, 1036
dieback,
attack density, 1363
Korean oak wilt, 1363
Mongolian oak, 1363
Platypus koryoensis, 1363
diel behavior,
adult distribution, 661
codling moth, 661
mating disruption, 661
vacuum, 661
dietary exposure assay,
Ephestia egg, 1613
Orius insidiosus, 1613
plant-incorporated protectants (PIPs), 1613
Tier-I toxicity assay, 1613
dietary self-selection,
behavior, 1285
insect rearing, 1285
nutrition, 1285
yellow mealworm, 1285
diffusion model,
Decapitating fly, 73
dispersal, 73
Solenopsis invicta, 73
disease emergence,
Hemiptera, 1345
syndrome 'basses richesses', 1345
vector, 1345
disease transmission,
alfalfa, 782
alfalfa leafcutting bees, 782
bees, 782
chalkbrood, 782
dispersal,
Decapitating fly, 73
diffusion model, 73
Solenopsis invicta, 73
disruptive selection,
frequency-dependent selection, 1530
mate choice, 1530
pheromone, 1530
distribution, 111, 281
carrier specificity, 281
Lucanidae, 281
negative binomial distribution, 111
nematode, 281
phoresy, 281
population ecology, 111
Sahlbergella singularis, 111
spatial dispersion model, 111
disturbance, 82, 1051
bioassay, 82
chemical attraction, 82
fire, 1051
insect, 1051
invertebrate, 1051
succession, 1051
Tribolium castaneum, 82
Dominican Republic,
Cuba, 989
Guadeloupe, 989
Martinique, 989
Trinidad, 989
double-decker traps,
Agrilus planipennis, 1239
Fraxinus, 1239
girdled trees, 1239
Manuka oil, 1239
Drosophila,
cytochrome P450, 1604
detoxification, 1604
mycophagy, 1604

E

E-64,
environmental risk assessment, 496
nontarget effects, 496
potassium arsenate, 496
tiered testing, 496
EAG,
annual bluegrass weevil, 412
GC-MS, 412

green leaf volatiles, 412
Listronotus maculicollis, 412
ecosystem functioning,
biodiversity, 8
biological control, 8
lady beetle, 8
predatory bugs, 8
ecosystem service,
biological control, 260
tritrophic interaction, 260
egg,
egg pod, 157
hatching, 157
oviposition, 157
soil moisture, 157
egg pod,
egg, 157
hatching, 157
oviposition, 157
soil moisture, 157
egg trap,
almonds, 706
Amyelois transitella, 706
monitoring, 706
Prunis dulcis, 706
Elatobium abietinum,
color response, 120
flight activity, 120
Sitka spruce, 120
sticky traps, 120
electronic tag,
entanglement, 94
harmonic radar, 94
insect tracking, 94
insect walking, 94
ELISA,
Bt protein expression, 1323
control efficacy, 1323
transgenic rice, 1323
elm leaf beetle,
cold hardiness, 1546
diapause, 1546
supercooling point, 1546
Xanthogaleruca luteola, 1546
elongation factor 1-α,
biotypes, 1317
cytochrome c, 1317
greenbug, 1317
host-plant resistance, 1317
EMEND,
forest management, 797
species composition, 797
variable retention, 797
vertical stratification, 797
entanglement,
Electronic tag, 94
harmonic radar, 94
insect tracking, 94
insect walking, 94
environmental risk assessment,
E-64, 496
nontarget effects, 496
potassium arsenate, 496
tiered testing, 496
enzyme inactivation,
Arrhenius plot, 462
intrinsic optimum temperature, 462
Sharpe-Schoolfield model, 462
tangent equation, 462
Eoreuma loftini (Dyar),
alternate hosts, 1036
Diatraea saccharalis (F.), 1036
Eoreuma loftini (Dyar), 1036
Mexican rice borer, 1036
sugarcane borer, 1036
EPG,
feeding behavior, 743
life table parameters, 743
transgenic GNA wheat, 743
wheat aphids, 743
Ephestia egg,
dietary exposure assay, 1613
Orius insidiosus, 1613
plant-incorporated protectants (PIPs), 1613
Tier-I toxicity assay, 1613
Epiphyas postvittana,
degree-days, 1370
invasive species, 1370
phenology, 1370
Eretmocerus eremicus,
Bemisia argentifolii, 36
Glomus etunicatum, 36
multi-trophic interactions, 36
silverleaf whitefly, 36
ergovaline,
alkaloids, 639
N-acetyl norloline, 639
Neotyphodium, 639
peramine, 639
Eurygaster integriceps,
geostatistics, 1253
SADIE, 1253
spatial distribution, 1253
Euschistus servus,
Nezara viridula, 303
Telenomus podisi, 303
Trissolcus basalis, 303
exocrine gland,
Brazilian pepper, 605
host specificity testing, 605
weed biological control, 605
extrafloral nectaries,
ants, 42
oriental fruit moth, 42
peach, 42
plant defense, 42
extreme temperatures,
climate change, 777
climate variation, 777
Pinus edulis, 777

F

family diversity,
family richness, 247
herbivorous, 247
landscape, 247
Vaccinium corymbosum, 247
family richness,
family diversity, 247
herbivorous, 247
landscape, 247
Vaccinium corymbosum, 247
farnesol,
β-farnesene, 420
Monitoring, 420
pear ester, 420
Z-3-hexenyl acetate, 420
fatty acids,
Hylastinus obscurus, 399
red clover, 399
Scolytidae, 399
Trifolium pratense, 399
fecundity,
cotton aphid, 431
lower developmental threshold, 431
temperature requirement, 431
thermal constant, 431
feeding behavior, 367, 484, 743
Bacillus thuringiensis, 484
cotton, 367
EPG, 743
Gossypium, 367
life table parameters, 743

Lygus, 367
refuge strategy, 484
simulation modeling, 484
transgenic crops, 484
transgenic GNA wheat, 743
western tarnished plant bug, 367
wheat aphids, 743
feeding preferences,
Abagrotis orbis, 1458
climbing cutworms, 1458
grapevine, 1458
host plant, 1458
female longevity,
Bt corn, 1585
head capsule width, 1585
male investment, 1585
stable isotopes, 1585
field validation,
developmental rate, 939
thermal threshold, 939
variable temperature, 939
Xylotrechus arvicola, 939
fire,
disturbance, 1051
insect, 1051
invertebrate, 1051
succession, 1051
flight activity,
color response, 120
Elatobium abietinum, 120
Sitka spruce, 120
sticky traps, 120
flight dispersal,
Mangifera indica, 873
thrips, 873
yellow sticky trap, 873
flight muscles,
coffee berry borer, 441
degeneration, 441
Hypothenemus hampei, 441
Florida hammocks,
insect herbivory, 1437
invasive weed, 1437
pastures, 1437
Solanum viarum, 1437
flower taps,
binomial sampling, 381
classification sampling, 381
operating characteristic, 381
visual inspection, 381
flowers,
conservation biocontrol, 1177
habitat management, 1177
native plants, 1177
natural enemies, 1177
food,
development, 1577
temperature, 1577
Typhlodromus athenas, 1577
food preference,
detection, 1276
foraging distance, 1276
model, 1276
surveillance, 1276
foraging,
body size ratio, 737
handling time, 737
temperature-size rule, 737
foraging distance,
detection, 1276
food preference, 1276
model, 1276
surveillance, 1276
forest damage,
cellular automata, 273
county pattern, 273
southern pine beetle, 273
forest fragmentation,
biomass, 239
carrion removal rates, 239
necrophagous beetles, 239
Scarabaeinae, 239
forest management,
EMEND, 797
species composition, 797
variable retention, 797
vertical stratification, 797
forest restoration,
genetic modification, 955
gypsy moth, 955
oxalate oxidase, 955
Formica neorufibarbis,
altitudinal range, 534
behavioral thermoregulation, 534
colonization, 534
nest site preference, 534
Formicidae,
Arachnida, 1523
Coleoptera, 1523
Micro-Hymenoptera, 1523
Winkler extractor, 1523
454 pyrosequencing,
Homalodisca vitripennis, 23
insect associated bacteria, 23
microbiome, 23
Frankliniella bispinosa,
inverse distance weighting, 893
kriging, 893
semivariogram, 893
Vaccinium corymbosum x *V. darrowi*, 893
frass,
cocoon, 126
Monodontomerus aeneus, 126
Osmia cornuta, 126
volatiles, 126
Fraxinus, 679, 1239
Agrilus planipennis, 1239
Buprestidae, 679
double-decker traps, 1239
girdled trees, 1239
invasive pest, 679
Manuka oil, 1239
sample unit, 679
sampling technique, 679
frequency-dependent selection,
disruptive selection, 1530
mate choice, 1530
pheromone, 1530
freshwater,
microsporidia, 289
nematodes, 289
parasitism, 289
streams, 289
fruit,
blueberry, 697
conservation biocontrol, 697
habitat management, 697
integrated pest management (IPM), 697
fruit set,
Cucurbitaceae, 614
fruit size, 614
oligolecty, 614
pollination efficiency, 614
fruit size,
Cucurbitaceae, 614
fruit set, 614
oligolecty, 614
pollination efficiency, 614

functional response,
attack rate, 835
handling time, 835
Homalotylus flaminius, 835
numerical response, 835
fungal pathogens,
mate choice, 1554
Metarhizium anisopliae, 1554
reproductive maturation, 1554
socially transmitted diseases, 1554
fungi-fungi interactions,
ant pests, 342
biological control, 342
induction of conidiation, 342
leaf-cutter ants, 342
fungicide,
Beauveria bassiana, 59
genetic improvement, 59
Metarhizium brunneum, 59
resistance, 59
fuscum,
Cerambycidae, 714
cinnamopterum, 714
contact pheromone, 714
Tetropium, 714

G

Galerucella,
biocontrol, 1074
colonization, 1074
Lythrum salicaria, 1074
matrix habitat, 1074
GC-MS,
annual bluegrass weevil, 412
EAG, 412
green leaf volatiles, 412
Listronotus maculicollis, 412
gene flow,
Coptotermes gestroi, 470
microsatellite markers, 470
population structure, 470
gene-for-gene interaction,
alternative host, 1303
landscape genetics, 1303
microsatellites, 1303
wheat, 1303
generalist insect predators,
consumption rate, 333
killing rate, 333
natural enemies, 333
spiders, 333
Generalized Linear Models,
additive diversity partitioning, 1081
IndVal, 1081
overgrazing, 1081
Scarabaeidae, 1081
genetic improvement,
Beauveria bassiana, 59
fungicide, 59
Metarhizium brunneum, 59
resistance, 59
genetic modification,
forest restoration, 955
gypsy moth, 955
oxalate oxidase, 955
genotype-temperature interactions,
body size, 1
development time, 1
temperature, 1
geographical distribution, 549, 1183
bark beetles, 549
Dendroctonus rhizophagus, 549
holly leaf-miner, 1183
host-parasitoid system, 1183
Ilex aquifolium, 1183
predictive models, 549
geostatistics, 173, 1253
apple, 173
Eurygaster integriceps, 1253
median polishing, 173
residuals, 173
SADIE, 1253
spatial distribution, 1253
Synanthedon scitula, 173
girdled trees, 357, 1239
Agrilus planipennis, 1239
double-decker traps, 1239
Fraxinus, 1239
green ash, 357
Invasive forest pest, 357
Manuka oil, 1239
methyl jasmonate, 357
phloem- and wood-boring insect, 357
GIS,
glassy-winged sharpshooter, 755
grape, 755
Homalodisca vitripennis, 755
invasive species, 755
glassy-winged sharpshooter, 755, 1192
GIS, 755
grape, 755, 1192
Homalodisca vitripennis, 755
invasive species, 755
Pierce's disease, 1192
susceptibility, 1192
transmission efficiency, 1192
Glenea cantor,
bark moisture content, 1487
colonization, 1487
host selection, 1487
oviposition attractant, 1487
Glomus etunicatum,
Bemisia argentifolii, 36
Eretmocerus eremicus, 36
multi-trophic interactions, 36
silverleaf whitefly, 36
glycogen,
Gryllus firmus, 1566
trehalose, 1566
triglyceride, 1566
wing dimorphism, 1566
goats,
catechin, 1572
cows, 1572
Musca domestica, 1572
golf courses,
introduction, 589
Ovavesicula popilliae, 589
Popillia japonica, 589
turfgrass, 589
Gossypium,
cotton, 367
feeding behavior, 367
Lygus, 367
western tarnished plant bug, 367
grape, 755, 1192
GIS, 755
glassy-winged sharpshooter, 755, 1192
Homalodisca vitripennis, 755
invasive species, 755
Pierce's disease, 1192
susceptibility, 1192
transmission efficiency, 1192
grape berry moth,
climate change, 505
greenhouse gas emissions, 505
phenology, 505

voltinism, 505
grapevine,
Abagrotis orbis, 1458
climbing cutworms, 1458
feeding preferences, 1458
host plant, 1458
grass aphid,
greenbug, 151
pea aphid, 151
Russian wheat aphid, 151
western wheat aphid, 151
green ash,
girdled trees, 357
Invasive forest pest, 357
methyl jasmonate, 357
phloem- and wood-boring insect, 357
green leaf volatiles,
annual bluegrass weevil, 412
EAG, 412
GC-MS, 412
Listronotus maculicollis, 412
green trap,
Agrilus planipennis, 904
kairomone, 904
lactone, 904
pheromone, 904
greenbug, 151, 1317
biotypes, 1317
cytochrome c, 1317
elongation factor 1-α, 1317
grass aphid, 151
host-plant resistance, 1317
pea aphid, 151
Russian wheat aphid, 151
western wheat aphid, 151
greenhouse gas emissions,
climate change, 505
grape berry moth, 505
phenology, 505
voltinism, 505
Gryllus firmus,
glycogen, 1566
trehalose, 1566
triglyceride, 1566
wing dimorphism, 1566
Guadeloupe,
Cuba, 989
Dominican Republic, 989
Martinique, 989
Trinidad, 989
gut bacteria,
16S rRNA-restriction fragment-length polymorphism, 1405
Camponotus japonicus, 1405
gut symbiont,
Burkholderia, 1102
Coreidae, 1102
crypts, 1102
Wolbachia, 1102
gypsy moth,
forest restoration, 955
genetic modification, 955
oxalate oxidase, 955

H

habitat management, 697, 1177
blueberry, 697
conservation biocontrol, 697, 1177
flowers, 1177
fruit, 697
integrated pest management (IPM), 697
native plants, 1177
natural enemies, 1177
habitat modification,
ants, 1019
invasive species, 1019
microclimate, 1019
niche expansion, 1019
habitat structure, 1378, 1397
active hunters, 1378
community ecology, 1397
landscaping, 1397
odorous house ant, 1397
predators, 1378
Tsuga canadensis, 1378
urban, 1397
web builders, 1378
handling time, 737, 835
attack rate, 835
body size ratio, 737
foraging, 737
functional response, 835
Homalotylus flaminius, 835
numerical response, 835
temperature-size rule, 737
harmonic radar,
Electronic tag, 94
entanglement, 94
insect tracking, 94
insect walking, 94
hatching,
egg, 157
egg pod, 157
oviposition, 157
soil moisture, 157
head capsule width,
Bt corn, 1585
female longevity, 1585
male investment, 1585
stable isotopes, 1585
head width,
Prunus avium, 1353
Prunus emarginata, 1353
Western cherry fruit fly, 1353
wing length, 1353
Heliothis subflexa,
neonate movement, 1478
nonhost oviposition, 1478
oviposition mistakes, 1478
Physalis, 1478
Hemiptera,
disease emergence, 1345
syndrome 'basses richesses', 1345
vector, 1345
hemlock woolly adelgid,
Adelges tsugae, 324
biological control, 324
Host range, 324
Laricobius osakensis, 324
herbivorous,
family diversity, 247
family richness, 247
landscape, 247
Vaccinium corymbosum, 247
Hermetia illucens,
Bacillus subtilis, 30
chicken manure, 30
waste management, 30
holly leaf-miner,
geographical distribution, 1183
host-parasitoid system, 1183
Ilex aquifolium, 1183
Homalodisca vitripennis, 23, 755
454 pyrosequencing, 23
GIS, 755
glassy-winged sharpshooter, 755
grape, 755
insect associated bacteria, 23
invasive species, 755

microbiome, 23
Homalotylus flaminius,
attack rate, 835
functional response, 835
handling time, 835
numerical response, 835
host,
body size, 931
Mediterranean fruit fly, 931
sterile insect technique, 931
temperature, 931
host finding,
host selection, 855
insect, 855
plant, 855
visual cues, 855
host finding behavior,
Adelges tsugae, 864
biological control, 864
huanglongbin, 1494
olfactory, 1494
plant–herbivore interactions, 1494
predator-prey interactions, 864
Tsuga spp., 864
visual, 1494
host plant,
Abagrotis orbis, 1458
climbing cutworms, 1458
feeding preferences, 1458
grapevine, 1458
host plant resistance,
Agrilus anxius, 648
biogeography, 648
coevolution, 648
wood-borers, 648
host quality,
biological control, 1123
competitive interaction, 1123
multiple release, 1123
population dynamics, 1123
host range,
Adelges tsugae, 324
biological control, 324
hemlock woolly adelgid, 324
Laricobius osakensis, 324
host parasitoid system,
geographical distribution, 1183
holly leaf-miner, 1183
Ilex aquifolium, 1183
host plant resistance,
biotypes, 1317
cytochrome c, 1317
elongation factor 1-α, 1317
greenbug, 1317
host selection, 855, 1487
bark moisture content, 1487
colonization, 1487
Glenea cantor, 1487
host finding, 855
insect, 855
oviposition attractant, 1487
plant, 855
visual cues, 855
host specificity,
Chrysomelidae, 1427
nontarget effects, 1427
prerelease impact, 1427
swallow-worts, 1427
host specificity testing,
Brazilian pepper, 605
exocrine gland, 605
weed biological control, 605
host utilization,
alatae, 350
intraguild competition, 350
Macrosiphum euphorbiae, 350
Myzus persicae, 350
HSP genes,
Bemisia tabaci, 132
cold hardiness, 132
rapid cold hardening, 132
Huanglongbing,
Asian citrus psyllid, 1503
biological control, 1503
Candidatus liberibacter asiaticus, 1503
host-finding behavior, 1494
life table, 1503
olfactory, 1494
plant–herbivore interactions, 1494
visual, 1494
Hylastinus obscurus,
fatty acids, 399
red clover, 399
Scolytidae, 399
Trifolium pratense, 399
Hypothenemus hampei,
coffee berry borer, 441
degeneration, 441
flight muscles, 441

I

Ilex aquifolium,
geographical distribution, 1183
holly leaf-miner, 1183
host-parasitoid system, 1183
imperfect mimicry,
Batesian mimicry, 1223
myrmecomorphy, 1223
Peckhamia, 1223
Salticidae, 1223
indirect interactions,
climate change, 824
community structure, 824
physiological ecology, 824
species interactions, 824
induction of conidiation,
ant pests, 342
biological control, 342
fungi-fungi interactions, 342
leaf-cutter ants, 342
IndVal,
additive diversity partitioning, 1081
Generalized Linear Models, 1081
overgrazing, 1081
Scarabaeidae, 1081
insect, 855, 1051
disturbance, 1051
fire, 1051
host finding, 855
host selection, 855
invertebrate, 1051
plant, 855
succession, 1051
visual cues, 855
insect associated bacteria,
454 pyrosequencing, 23
Homalodisca vitripennis, 23
microbiome, 23
insect conservation,
Curculionidae, 1388
phytophagy, 1388
plant-insect interaction, 1388
insect herbivory,
Florida hammocks, 1437
invasive weed, 1437
pastures, 1437
Solanum viarum, 1437
insect host-parasitoid,
biosafety, 1331
leafy gene, 1331

np-ENDITALII selectable marker gene, 1331
risk assessment, 1331
insect-microbe interactions,
midgut microbiota, 1111
Rothschildia lebeau, 1111
tropical caterpillar, 1111
Área de Conservación Guanacaste, 1111
insect-plant interactions,
nitrogen, 66
phosphorus, 66
potassium, 66
saltcedar, 66
insect rearing,
behavior, 1285
dietary self-selection, 1285
nutrition, 1285
yellow mealworm, 1285
insect resistance management, 477, 964
pest behavior, 964
refuge, 964
refugia, 477
resistance evolution, 477
seed mixture, 964
spatial dynamics, 964
insect tracking,
Electronic tag, 94
entanglement, 94
harmonic radar, 94
insect walking, 94
insect walking,
Electronic tag, 94
entanglement, 94
harmonic radar, 94
insect tracking, 94
insecticidal crystalline proteins,
Bacillus thuringiensis subsp. *israelensis*, 1417
diatoms, 1417
seston, 1417
Simulium vittatum, 1417
insects,
copper binding proteins, 167
nano-RP-HPLC–ESI–MS, 167
integrated pest management (IPM),
blueberry, 697
conservation biocontrol, 697
fruit, 697
habitat management, 697
pecan, 889
pecan nut casebearer, 889
sex pheromone, 889
interspecific interaction,
alien species, 140
Bemisia tabaci, 140
co-occurrence, 140
Trialeurodes vaporariorum, 140
intraguild competition,
alatae, 350
host utilization, 350
Macrosiphum euphorbiae, 350
Myzus persicae, 350
intraguild predation,
Biological control, 103
cannibalism, 103
Coccinellidae, 103
larval tracks, 103
intrinsic optimum temperature,
Arrhenius plot, 462
enzyme inactivation, 462
Sharpe-Schoolfield model, 462
tangent equation, 462
introduction,
golf courses, 589
Ovavesicula popilliae, 589
Popillia japonica, 589
turfgrass, 589
inundation,
larval gas exchange, 1295
plastron, 1295
salt water, 1295
tidal activity, 1295
invasive forest pest,
girdled trees, 357
green ash, 357
methyl jasmonate, 357
phloem- and wood-boring insect, 357
invasive fruit fly,
Bactrocera dorsalis complex, 844
polyphagous, 844
quarantine pest, 844
Tephritidae, 844
invasive insect,
conifer pest, 1266
native parasitoids, 1266
pine mortality, 1266
invasive pest,
Buprestidae, 679
Fraxinus, 679
sample unit, 679
sampling technique, 679
invasive species, 231, 755, 1019, 1370
ants, 1019
degree-days, 1370
Epiphyas postvittana, 1370
GIS, 755
glassy-winged sharpshooter, 755
grape, 755
habitat modification, 1019
Homalodisca vitripennis, 755
microclimate, 1019
niche expansion, 1019
phenology, 1370
population monitoring, 231
seasonality, 231
species composition, 231
stink bug, 231
invasive weed,
Florida hammocks, 1437
insect herbivory, 1437
pastures, 1437
Solanum viarum, 1437
inventory,
bees, 374
bowl trapping, 374
netting, 374
sampling, 374
inverse distance weighting,
Frankliniella bispinosa, 893
kriging, 893
semivariogram, 893
Vaccinium corymbosum x *V. darrowi*, 893
invertebrate,
disturbance, 1051
fire, 1051
insect, 1051
succession, 1051
IPM,
biological control, 1093
Carabidae, 1093
Cyclotrachelus sodalis, 1093
omnivory, 1093
Italian ryegrass,
agroecosystem, 1027
apparent competition, 1027
plant introduction, 1027

K

kairomone,
Agrilus planipennis, 904
green trap, 904
lactone, 904
pheromone, 904
killing rate,
consumption rate, 333
generalist insect predators, 333
natural enemies, 333
spiders, 333
Konza Prairie Biological Station,
life-history, 979
management, 979
Orthoptera, 979
reproduction, 979
Korean oak wilt,
attack density, 1363
dieback, 1363
Mongolian oak, 1363
Platypus koryoensis, 1363
kriging,
Frankliniella bispinosa, 893
inverse distance weighting, 893
semivariogram, 893
Vaccinium corymbosum x *V. darrowi*, 893

L

lactone,
Agrilus planipennis, 904
green trap, 904
kairomone, 904
pheromone, 904
lady beetle,
biodiversity, 8
biological control, 8
ecosystem functioning, 8
predatory bugs, 8
landscape,
family diversity, 247
family richness, 247
herbivorous, 247
Vaccinium corymbosum, 247
landscape ecology,
agro-ecosystem, 621
Andrenidae, 621
apple orchards, 621
native bees, 621
landscape genetics,
alternative host, 1303
gene-for-gene interaction, 1303
microsatellites, 1303
wheat, 1303
landscaping,
community ecology, 1397
habitat structure, 1397
odorous house ant, 1397
urban, 1397
Laricobius osakensis,
Adelges tsugae, 324
biological control, 324
hemlock woolly adelgid, 324
Host range, 324
larval development, 184, 633
alternate hosts, 633
Diabrotica undecimpunctata undecimpunctata, 633
population dynamics, 184
Stomoxys calcitrans, 184
larval gas exchange,
inundation, 1295
plastron, 1295
salt water, 1295
tidal activity, 1295
larval parasitoid,
Bactrocera oleae (Rossi), 315
Olea europaea L., 315
larval tracks,
Biological control, 103
cannibalism, 103
Coccinellidae, 103
intraguild predation, 103
leaf-cutter ants,
ant pests, 342
biological control, 342
fungi-fungi interactions, 342
induction of conidiation, 342
leaf-galls,
biological control, 1448
Lygodium microphyllum, 1448
leafy gene,
biosafety, 1331
insect host-parasitoid, 1331
*np-ENDITAL*II selectable marker gene, 1331
risk assessment, 1331
Leucopis argenticollis,
Adelges tsugae, 1410
biological control, 1410
Leucopis piniperda, 1410
prey suitability, 1410
Leucopis piniperda,
Adelges tsugae, 1410
biological control, 1410
Leucopis argenticollis, 1410
prey suitability, 1410
life history,
Konza Prairie Biological Station, 979
management, 979
Orthoptera, 979
reproduction, 979
life table, 597, 1503
Asian citrus psyllid, 1503
biological control, 1503
Candidatus liberibacter asiaticus, 1503
development, 597
Huanglongbing, 1503
Phytoseiidae, 597
pollen., 597
Typhlodromus athenas, 597
life table parameters,
EPG, 743
feeding behavior, 743
transgenic GNA wheat, 743
wheat aphids, 743
lipids,
alfalfa leafcutting bee, 917
Megachile rotundata, 917
pollinators, 917
thermal biology, 917
Liriomyza huidobrensis,
biological control, 217
Liriomyza sativae, 217
parasitoids, 217
Liriomyza sativae,
biological control, 217
Liriomyza huidobrensis, 217
parasitoids, 217
Listronotus maculicollis,
annual bluegrass weevil, 412
EAG, 412
GC-MS, 412
green leaf volatiles, 412
lower developmental threshold,
cotton aphid, 431
fecundity, 431
temperature requirement, 431
thermal constant, 431

Lucanidae,
carrier specificity, 281
distribution, 281
nematode, 281
phoresy, 281
Lygodium microphyllum,
biological control, 1448
leaf-galls, 1448
Lygus hesperus,
diapause, 455
rearing, 455
western tarnished plant bug, 455
Lygus,
cotton, 367
feeding behavior, 367
Gossypium, 367
western tarnished plant bug, 367
Lythrum salicaria,
biocontrol, 1074
colonization, 1074
Galerucella, 1074
matrix habitat, 1074

M

Macrosiphum euphorbiae,
alatae, 350
host utilization, 350
intraguild competition, 350
Myzus persicae, 350
male investment,
Bt corn, 1585
female longevity, 1585
head capsule width, 1585
stable isotopes, 1585
management,
Konza Prairie Biological Station, 979
life-history, 979
Orthoptera, 979
reproduction, 979
Mangifera indica,
flight dispersal, 873
thrips, 873
yellow sticky trap, 873
Manuka oil,
Agrilus planipennis, 1239
double-decker traps, 1239
Fraxinus, 1239
girdled trees, 1239
Martinique,
Cuba, 989
Dominican Republic, 989
Guadeloupe, 989
Trinidad, 989
mass trapping,
computer simulation in three dimensions, 1210
mating disruption, 1210
mean flight height, 1210
movement algorithms, 1210
mate choice, 1530, 1554
disruptive selection, 1530
frequency-dependent selection, 1530
fungal pathogens, 1554
Metarhizium anisopliae, 1554
pheromone, 1530
reproductive maturation, 1554
socially transmitted diseases, 1554
mating disruption, 661, 1210
adult distribution, 661
codling moth, 661
computer simulation in three dimensions, 1210
diel behavior, 661
mass trapping, 1210
mean flight height, 1210
movement algorithms, 1210
vacuum, 661
matrix habitat,
biocontrol, 1074
colonization, 1074
Galerucella, 1074
Lythrum salicaria, 1074
mean flight height,
computer simulation in three dimensions, 1210
mass trapping, 1210
mating disruption, 1210
movement algorithms, 1210
median polishing,
apple, 173
geostatistics, 173
residuals, 173
Synanthedon scitula, 173
Mediterranean fruit fly,
body size, 931
host, 931
sterile insect technique, 931
temperature, 931
Megachile rotundata,
alfalfa leafcutting bee, 917
lipids, 917
pollinators, 917
thermal biology, 917
Membracidae,
Aphrophoridae, 1131
Cicadellidae, 1131
Clastopteridae, 1131
Xylella fastidiosa, 1131
Metarhizium anisopliae,
fungal pathogens, 1554
mate choice, 1554
reproductive maturation, 1554
socially transmitted diseases, 1554
Metarhizium brunneum,
Beauveria bassiana, 59
fungicide, 59
genetic improvement, 59
resistance, 59
methyl jasmonate,
girdled trees, 357
green ash, 357
Invasive forest pest, 357
phloem- and wood-boring insect, 357
Mexican rice borer,
alternate hosts, 1036
Diatraea saccharalis (F.), 1036
Eoreuma loftini (Dyar), 1036
sugarcane borer, 1036
Micro-Hymenoptera,
Arachnida, 1523
Coleoptera, 1523
Formicidae, 1523
Winkler extractor, 1523
microbiome,
454 pyrosequencing, 23
Homalodisca vitripennis, 23
insect associated bacteria, 23
microclimate,
ants, 1019
habitat modification, 1019
invasive species, 1019
niche expansion, 1019
microsatellite markers,
Coptotermes gestroi, 470
gene flow, 470
population structure, 470

microsatellites,
alternative host, 1303
gene-for-gene interaction, 1303
landscape genetics, 1303
wheat, 1303
microsporidia,
freshwater, 289
nematodes, 289
parasitism, 289
streams, 289
midgut microbiota,
insect-microbe interactions, 1111
Rothschildia lebeau, 1111
tropical caterpillar, 1111
Área de Conservación Guanacaste, 1111
migration, 449, 516
Chrysoperla sinica, 449
Mythimna separata, 516
ovarian development, 449
regulation, 516
tethered-flight, 449
MLST,
canola pest, 818
multilocus sequence typing, 818
symbiont, 818
Wolbachia, 818
model,
detection, 1276
food preference, 1276
foraging distance, 1276
surveillance, 1276
moisture,
Coptotermes formosanus, 1232
mortality, 1232
starvation, 1232
Mongolian oak,
attack density, 1363
dieback, 1363
Korean oak wilt, 1363
Platypus koryoensis, 1363
monitoring,
almonds, 706
Amyelois transitella, 706
beta-farnesene, 420
egg trap, 706
farnesol, 420
pear ester, 420
Prunis dulcis, 706
Z-3-hexenyl acetate, 420
Monodontomerus aeneus,
cocoon, 126
frass, 126
Osmia cornuta, 126
volatiles, 126
mortality,
Coptotermes formosanus, 1232
moisture, 1232
starvation, 1232
moth diversity,
agro-ecosystems, 560
organic farming, 560
plant diversity, 560
woody hedgerow, 560
movement algorithms,
computer simulation in three dimensions, 1210
mass trapping, 1210
mating disruption, 1210
mean flight height, 1210
multi-trophic interactions,
Bemisia argentifolii, 36
Eretmocerus eremicus, 36
Glomus etunicatum, 36
silverleaf whitefly, 36
multilocus sequence typing,
canola pest, 818
MLST, 818
symbiont, 818
Wolbachia, 818
multiple release,
biological control, 1123
competitive interaction, 1123
host quality, 1123
population dynamics, 1123
multiplex PCR,
pest identification, 1595
Planococcus, 1595
Pseudococcus, 1595
vineyards, 1595
Musca domestica, 52, 1572
Bacillus thuringiensis variety *israelensis*, 52
biological control, 52
catechin, 1572
cows, 1572
goats, 1572
poultry manure, 52
Muscidifurax raptorellus,
parasitism, 88
Spalangia cameroni, 88
Spalangia endius, 88
Stomoxys calcitrans, 88
mycophagy,
cytochrome P450, 1604
detoxification, 1604
Drosophila, 1604
myrmecomorphy,
Batesian mimicry, 1223
imperfect mimicry, 1223
Peckhamia, 1223
Salticidae, 1223
Mythimna separata,
migration, 516
regulation, 516
Myzus persicae, 350, 391
alatae, 350
Aphis gossypii, 391
barrier crop, 391
host utilization, 350
intraguild competition, 350
Macrosiphum euphorbiae, 350
nonpersistent virus, 391
virus sink, 391

N

N-acetyl norloline,
alkaloids, 639
ergovaline, 639
Neotyphodium, 639
peramine, 639
nano-RP-HPLC–ESI–MS,
copper binding proteins, 167
insects, 167
native bees,
agro-ecosystem, 621
Andrenidae, 621
apple orchards, 621
landscape ecology, 621
native parasitoids,
conifer pest, 1266
invasive insect, 1266
pine mortality, 1266
native plants,
conservation biocontrol, 1177
flowers, 1177
habitat management, 1177
natural enemies, 1177
native sunflowers,
Cylindrocopturus adspersus, 15
native versus cultivated, 15
parasitism, 15

native versus cultivated,
Cylindrocopturus adspersus, 15
native sunflowers, 15
parasitism, 15
natural enemies, 333, 1007, 1177
conservation biocontrol, 1177
consumption rate, 333
defenses, 1007
flowers, 1177
generalist insect predators, 333
habitat management, 1177
killing rate, 333
native plants, 1177
nutritional quality, 1007
outbreak, 1007
Scolytinae, 1007
spiders, 333
natural refuge,
nonrice host plants, 749
resistance management, 749
trap crops, 749
necrophagous beetles,
biomass, 239
carrion removal rates, 239
forest fragmentation, 239
Scarabaeinae, 239
negative binomial distribution,
distribution, 111
population ecology, 111
Sahlbergella singularis, 111
spatial dispersion model, 111
nematode,
carrier specificity, 281
distribution, 281
freshwater, 289
Lucanidae, 281
microsporidia, 289
parasitism, 289
phoresy, 281
streams, 289
neonate movement,
Heliothis subflexa, 1478
nonhost oviposition, 1478
oviposition mistakes, 1478
Physalis, 1478
Neotyphodium,
alkaloids, 639
ergovaline, 639
N-acetyl norloline, 639
peramine, 639
nest site preference,
altitudinal range, 534
behavioral thermoregulation, 534
colonization, 534
Formica neorufibarbis, 534
netting,
bees, 374
bowl trapping, 374
inventory, 374
sampling, 374
Nezara viridula,
Euschistus servus, 303
Telenomus podisi, 303
Trissolcus basalis, 303
niche expansion,
ants, 1019
habitat modification, 1019
invasive species, 1019
microclimate, 1019
Nicrophorus,
antimicrobial peptides, 669
competition, 669
parental care, 669
secretions, 669
nitrogen,
insect–plant interactions, 66
phosphorus, 66
potassium, 66
saltcedar, 66
nonconsumptive effects,
agroforestry, 581
Azteca instabilis, 581
Pseudacteon, 581
trait-meditated interactions, 581
nonhost oviposition,
Heliothis subflexa, 1478
neonate movement, 1478
oviposition mistakes, 1478
Physalis, 1478
nonpersistent virus,
Aphis gossypii, 391
barrier crop, 391
Myzus persicae, 391
virus sink, 391
nonrice host plants,
natural refuge, 749
resistance management, 749
trap crops, 749
nontarget effects, 496, 1427
Chrysomelidae, 1427
E-64, 496
environmental risk assessment, 496
host specificity, 1427
potassium arsenate, 496
prerelease impact, 1427
swallow-worts, 1427
tiered testing, 496
np-ENDITALII selectable marker gene,
biosafety, 1331
insect host-parasitoid, 1331
leafy gene, 1331
risk assessment, 1331
numerical response,
attack rate, 835
functional response, 835
handling time, 835
Homalotylus flaminius, 835
nutrition,
behavior, 1285
dietary self-selection, 1285
insect rearing, 1285
yellow mealworm, 1285
nutritional quality,
defenses, 1007
natural enemies, 1007
outbreak, 1007
Scolytinae, 1007

O

odorous house ant,
community ecology, 1397
habitat structure, 1397
landscaping, 1397
urban, 1397
Olea europaea L.,
Bactrocera oleae (Rossi), 315
Larval parasitoid, 315
olfactory,
host-finding behavior, 1494
huanglongbin, 1494
plant–herbivore interactions, 1494
visual, 1494
oligolecty,
Cucurbitaceae, 614
fruit set, 614
fruit size, 614
pollination efficiency, 614

omnivory,
biological control, 1093
Carabidae, 1093
Cyclotrachelus sodalis, 1093
IPM, 1093
operating characteristic,
binomial sampling, 381
classification sampling, 381
flower taps, 381
visual inspection, 381
organic crucifer production,
pest management, 1471
trap crop, 1471
organic farming,
agro-ecosystems, 560
moth diversity, 560
plant diversity, 560
woody hedgerow, 560
oriental fruit moth,
ants, 42
extrafloral nectaries, 42
peach, 42
plant defense, 42
Orius insidiosus,
dietary exposure assay, 1613
Ephestia egg, 1613
plant-incorporated protectants (PIPs), 1613
Tier-I toxicity assay, 1613
Orthoptera, 541, 979
Acrididae, 541
bottom-up forces, 541
Konza Prairie Biological Station, 979
life-history, 979
management, 979
reproduction, 979
resource limitation, 541
subarctic, 541
Osmia cornuta,
cocoon, 126
frass, 126
Monodontomerus aeneus, 126
volatiles, 126
outbreak,
defenses, 1007
natural enemies, 1007
nutritional quality, 1007
Scolytinae, 1007
ovarian development,
Chrysoperla sinica, 449
migration, 449
tethered-flight, 449
Ovavesicula popilliae,
golf courses, 589
introduction, 589
Popillia japonica, 589
turfgrass, 589
overgrazing,
additive diversity partitioning, 1081
Generalized Linear Models, 1081
IndVal, 1081
Scarabaeidae, 1081
oviposition,
egg, 157
egg pod, 157
hatching, 157
soil moisture, 157
oviposition attractant,
bark moisture content, 1487
colonization, 1487
Glenea cantor, 1487
host selection, 1487
oviposition mistakes,
Heliothis subflexa, 1478
neonate movement, 1478
nonhost oviposition, 1478
Physalis, 1478
oxalate oxidase,
forest restoration, 955
genetic modification, 955
gypsy moth, 955

P

palatable forage hypothesis,
Amazon, 770
Atta sexdens, 770
Colombia, 770
demography, 770
para-ethylacetophenone,
Anastrepha, 727
Bactrocera, 727
Ceratitis, 727
Diachasmimorpha longicaudata, 727
parasitism, 15, 88, 289
Cylindrocopturus adspersus, 15
freshwater, 289
microsporidia, 289
Muscidifurax raptorellus, 88
native sunflowers, 15
native versus cultivated, 15
nematodes, 289
Spalangia cameroni, 88
Spalangia endius, 88
Stomoxys calcitrans, 88
streams, 289
parasitoids,
biological control, 217
Liriomyza huidobrensis, 217
Liriomyza sativae, 217
parental care,
antimicrobial peptides, 669
competition, 669
Nicrophorus, 669
secretions, 669
pastures, 999, 1437
ant mosaic, 999
ants, 999
Florida hammocks, 1437
insect herbivory, 1437
invasive weed, 1437
scramble competition, 999
Solanum viarum, 1437
spatial pattern, 999
path analysis,
aphids, 570
coexistence, 570
population regulation, 570
pea aphid,
grass aphid, 151
greenbug, 151
Russian wheat aphid, 151
western wheat aphid, 151
peach,
ants, 42
extrafloral nectaries, 42
oriental fruit moth, 42
plant defense, 42
peanut,
brown stink bug, 1159
cotton, 1159
Southern green stink bug, 1159
stink bug distribution, 1159
pear ester,
β-farnesene, 420
farnesol, 420
Monitoring, 420
(*Z*)-3-hexenyl acetate, 420
pecan,
integrated pest management (IPM), 889
pecan nut casebearer, 889

sex pheromone, 889
pecan nut casebearer,
integrated pest management (IPM), 889
pecan, 889
sex pheromone, 889
Peckhamia,
Batesian mimicry, 1223
imperfect mimicry, 1223
myrmecomorphy, 1223
Salticidae, 1223
Peponapis pruinosa,
Bombus impatiens, 1144
Cucurbita spp., 1144
pollination, 1144
peramine,
alkaloids, 639
ergovaline, 639
N-acetyl norloline, 639
Neotyphodium, 639
pest behavior,
insect resistance management, 964
refuge, 964
seed mixture, 964
spatial dynamics, 964
pest identification,
multiplex PCR, 1595
Planococcus, 1595
Pseudococcus, 1595
vineyards, 1595
pest management, 689, 1471
aggregation, 689
autocorrelation, 689
organic crucifer production, 1471
Semivariogram, 689
Taylor's power law, 689
trap crop, 1471
phenology, 505, 1370, 1511
climate change, 505
degree-days, 1370
Epiphyas postvittana, 1370
grape berry moth, 505
greenhouse gas emissions, 505
invasive species, 1370
population, 1511
spatial distribution, 1511
Tortricidae, 1511
volatiles, 1511
voltinism, 505
pheromone, 654, 904, 1530
Agrilus planipennis, 904
disruptive selection, 1530
frequency-dependent selection, 1530
green trap, 904
kairomone, 904
lactone, 904
mate choice, 1530
precision pest management, 654
spatial, 654
Striacosta albicosta, 654
western bean cutworm, 654
phloem- and wood-boring insect,
girdled trees, 357
green ash, 357
Invasive forest pest, 357
methyl jasmonate, 357
phoresy,
carrier specificity, 281
distribution, 281
Lucanidae, 281
nematode, 281
phosphorus,
insect–plant interactions, 66
nitrogen, 66
potassium, 66
saltcedar, 66
Physalis,
Heliothis subflexa, 1478
neonate movement, 1478
nonhost oviposition, 1478
oviposition mistakes, 1478
physiological ecology,
climate change, 824
community structure, 824
indirect interactions, 824
species interactions, 824
phytophagy,
Curculionidae, 1388
insect conservation, 1388
plant-insect interaction, 1388
Phytoseiidae,
development, 597
life table, 597
pollen., 597
Typhlodromus athenas, 597
Pierce's disease,
glassy-winged sharpshooter, 1192
grape, 1192
Pierce's disease, 1192
susceptibility, 1192
transmission efficiency, 1192
pine mortality,
conifer pest, 1266
invasive insect, 1266
native parasitoids, 1266
Pinus edulis,
climate change, 777
climate variation, 777
extreme temperatures, 777
Planococcus,
multiplex PCR, 1595
pest identification, 1595
Pseudococcus, 1595
vineyards, 1595
plant,
host finding, 855
host selection, 855
insect, 855
visual cues, 855
plant defense,
ants, 42
extrafloral nectaries, 42
oriental fruit moth, 42
peach, 42
plant diversity,
agro-ecosystems, 560
moth diversity, 560
organic farming, 560
woody hedgerow, 560
plant introduction,
agroecosystem, 1027
apparent competition, 1027
Italian ryegrass, 1027
plant-incorporated protectants (PIPs),
dietary exposure assay, 1613
Ephestia egg, 1613
Orius insidiosus, 1613
Tier-I toxicity assay, 1613
plant-insect interaction,
behavior, 880
Curculionidae, 1388
insect conservation, 1388
phytophagy, 1388
roots, 880
Vitacea polistiformis, 880
plant–herbivore interactions,
host-finding behavior, 1494
huanglongbin, 1494
olfactory, 1494

visual, 1494
plastron,
inundation, 1295
larval gas exchange, 1295
salt water, 1295
tidal activity, 1295
Platypus koryoensis,
attack density, 1363
dieback, 1363
Korean oak wilt, 1363
Mongolian oak, 1363
pollen.,
development, 597
life table, 597
Phytoseiidae, 597
Typhlodromus athenas, 597
pollination,
Bombus impatiens, 1144
Cucurbita spp., 1144
Peponapis pruinosa, 1144
pollination efficiency,
Cucurbitaceae, 614
fruit set, 614
fruit size, 614
oligolecty, 614
pollinators,
alfalfa leafcutting bee, 917
lipids, 917
Megachile rotundata, 917
thermal biology, 917
polyphagous,
Bactrocera dorsalis complex, 844
invasive fruit fly, 844
quarantine pest, 844
Tephritidae, 844
Popillia japonica,
golf courses, 589
introduction, 589
Ovavesicula popilliae, 589
turfgrass, 589
population,
phenology, 1511
spatial distribution, 1511
Tortricidae, 1511
volatiles, 1511
population dynamics, 184, 1123
biological control, 1123
competitive interaction, 1123
host quality, 1123
larval development, 184
multiple release, 1123
Stomoxys calcitrans, 184
population ecology, 111, 194
Allium cepa, 194
distribution, 111
negative binomial distribution, 111
Sahlbergella singularis, 111
spatial dispersion model, 111
Thrips tabaci, 194
Tospovirus, 194
population growth,
developmental thresholds, 788
stored grain, 788
stored products, 788
population monitoring,
invasive species, 231
seasonality, 231
species composition, 231
stink bug, 231
population regulation,
aphids, 570
coexistence, 570
path analysis, 570
population structure,
Coptotermes gestroi, 470
gene flow, 470
microsatellite markers, 470
potassium,
insect–plant interactions, 66
nitrogen, 66
phosphorus, 66
saltcedar, 66
potassium arsenate,
E-64, 496
environmental risk assessment, 496
nontarget effects, 496
tiered testing, 496
poultry manure,
Bacillus thuringiensis variety *israelensis*, 52
biological control, 52
Musca domestica, 52
precision pest management,
pheromone, 654
spatial, 654
Striacosta albicosta, 654
western bean cutworm, 654
predation,
developmental time, 948
spined soldier bug, 948
yellow margined leaf beetle, 948
predator,
active hunters, 1378
aphids, 1541
habitat structure, 1378
pumpkin, 1541
rearing, 1541
thermal thresholds, 1541
Tsuga canadensis, 1378
web builders, 1378
predator-prey interactions,
Adelges tsugae, 864
biological control, 864
host finding behavior, 864
Tsuga spp., 864
predatory bugs,
biodiversity, 8
biological control, 8
ecosystem functioning, 8
lady beetle, 8
predictive models,
bark beetles, 549
Dendroctonus rhizophagus, 549
geographical distribution, 549
prerelease impact,
Chrysomelidae, 1427
host specificity, 1427
nontarget effects, 1427
swallow-worts, 1427
prey suitability,
Adelges tsugae, 1410
biological control, 1410
Leucopis argenticollis, 1410
Leucopis piniperda, 1410
Prunis dulcis,
almonds, 706
Amyelois transitella, 706
egg trap, 706
monitoring, 706
Prunus angustifolia,
Prunus persica, 1465
Prunus serotina, 1465
Synanthedon exitiosa, 1465
Synanthedon pictipes, 1465
Prunus avium,
head width, 1353
Prunus emarginata, 1353
Western cherry fruit fly, 1353
wing length, 1353

Prunus emarginata,
head width, 1353
Prunus avium, 1353
Western cherry fruit fly, 1353
wing length, 1353
Prunus persica,
Prunus angustifolia, 1465
Prunus serotina, 1465
Synanthedon exitiosa, 1465
Synanthedon pictipes, 1465
Prunus serotina,
Prunus angustifolia, 1465
Prunus persica, 1465
Synanthedon exitiosa, 1465
Synanthedon pictipes, 1465
Pseudacteon,
agroforestry, 581
Azteca instabilis, 581
nonconsumptive effects, 581
trait-meditated interactions, 581
Pseudococcus,
multiplex PCR, 1595
pest identification, 1595
Planococcus, 1595
vineyards, 1595
pumpkin,
aphids, 1541
predator, 1541
rearing, 1541
thermal thresholds, 1541

Q

quarantine pest,
Bactrocera dorsalis complex, 844
invasive fruit fly, 844
polyphagous, 844
Tephritidae, 844

R

rapid cold hardening,
Bemisia tabaci, 132
cold hardiness, 132
HSP genes, 132
rearing, 455, 1541
aphids, 1541
diapause, 455
Lygus hesperus, 455
predator, 1541
pumpkin, 1541
thermal thresholds, 1541
western tarnished plant bug, 455
red clover,
fatty acids, 399
Hylastinus obscurus, 399
Scolytidae, 399
Trifolium pratense, 399
refuge,
insect resistance management, 964
pest behavior, 964
seed mixture, 964
spatial dynamics, 964
refuge strategy,
Bacillus thuringiensis, 484
feeding behavior, 484
simulation modeling, 484
transgenic crops, 484
refugia,
insect resistance management, 477
resistance evolution, 477
regulation,
migration, 516
Mythimna separata, 516
reproduction,
Konza Prairie Biological Station, 979
life-history, 979
management, 979
Orthoptera, 979
reproductive maturation,
fungal pathogens, 1554
mate choice, 1554
Metarhizium anisopliae, 1554
socially transmitted diseases, 1554
residuals,
apple, 173
geostatistics, 173
median polishing, 173
Synanthedon scitula, 173
resistance,
Beauveria bassiana, 59
fungicide, 59
genetic improvement, 59
Metarhizium brunneum, 59
resistance evolution,
insect resistance management, 477
refugia, 477
resistance management,
natural refuge, 749
nonrice host plants, 749
trap crops, 749
resistance to crucifer flea beetles,
Arabidopsis, 1622
β-ionone, 1622
carotenoid cleavage dioxygenase one over-expression, 1622
resource limitation,
Acrididae, 541
bottom-up forces, 541
Orthoptera, 541
subarctic, 541
risk assessment,
biosafety, 1331
insect host-parasitoid, 1331
leafy gene, 1331
np-ENDITALII selectable marker gene, 1331
roots,
behavior, 880
plant-insect interactions, 880
Vitacea polistiformis, 880
Rothschildia lebeau,
insect-microbe interactions, 1111
midgut microbiota, 1111
tropical caterpillar, 1111
Área de Conservación Guanacaste, 1111
Russian wheat aphid,
grass aphid, 151
greenbug, 151
pea aphid, 151
western wheat aphid, 151

S

SADIE,
Eurygaster integriceps, 1253
geostatistics, 1253
spatial distribution, 1253
Sahlbergella singularis,
distribution, 111
negative binomial distribution, 111
population ecology, 111
spatial dispersion model, 111
salt water,
inundation, 1295
larval gas exchange, 1295

plastron, 1295
tidal activity, 1295
saltcedar,
insect–plant interactions, 66
nitrogen, 66
phosphorus, 66
potassium, 66
Salticidae,
Batesian mimicry, 1223
imperfect mimicry, 1223
myrmecomorphy, 1223
Peckhamia, 1223
sample unit,
Buprestidae, 679
Fraxinus, 679
invasive pest, 679
sampling technique, 679
sampling,
bees, 374
bowl trapping, 374
inventory, 374
netting, 374
sampling technique,
Buprestidae, 679
Fraxinus, 679
invasive pest, 679
sample unit, 679
Scarabaeidae,
additive diversity partitioning, 1081
Generalized Linear Models, 1081
IndVal, 1081
overgrazing, 1081
Scarabaeinae,
biomass, 239
carrion removal rates, 239
forest fragmentation, 239
necrophagous beetles, 239
Scolytidae,
fatty acids, 399
Hylastinus obscurus, 399
red clover, 399
Trifolium pratense, 399
Scolytinae,
defenses, 1007
natural enemies, 1007
nutritional quality, 1007
outbreak, 1007
scramble competition,
ant mosaic, 999
ants, 999
pastures, 999
spatial pattern, 999
seasonality,
invasive species, 231
population monitoring, 231
species composition, 231
stink bug, 231
secondary symbionts,
Bemisia tabaci, 809
Tunisia, 809
secretions,
antimicrobial peptides, 669
competition, 669
Nicrophorus, 669
parental care, 669
seed mixture,
insect resistance management, 964
pest behavior, 964
refuge, 964
spatial dynamics, 964
self-organization,
ant mosaics, 1067
spatial patterns, 1067
semivariogram,
aggregation, 689
autocorrelation, 689
Frankliniella bispinosa, 893
inverse distance weighting, 893
kriging, 893
pest management, 689
Taylor's power law, 689
Vaccinium corymbosum x *V. darrowi*, 893
seston,
Bacillus thuringiensis subsp. *israelensis*, 1417
diatoms, 1417
insecticidal crystalline proteins, 1417
Simulium vittatum, 1417
sex pheromone,
integrated pest management (IPM), 889
pecan, 889
pecan nut casebearer, 889
Sharpe-Schoolfield model,
Arrhenius plot, 462
enzyme inactivation, 462
intrinsic optimum temperature, 462
tangent equation, 462
silverleaf whitefly,
Bemisia argentifolii, 36
Eretmocerus eremicus, 36
Glomus etunicatum, 36
multi-trophic interactions, 36
simulation modeling,
Bacillus thuringiensis, 484
feeding behavior, 484
refuge strategy, 484
transgenic crops, 484
Simulium vittatum,
Bacillus thuringiensis subsp. *israelensis*, 1417
diatoms, 1417
insecticidal crystalline proteins, 1417
seston, 1417
Sitka spruce,
color response, 120
Elatobium abietinum, 120
flight activity, 120
sticky traps, 120
16S rRNA-restriction fragment-length polymorphism,
Camponotus japonicus, 1405
gut bacteria, 1405
socially transmitted diseases,
fungal pathogens, 1554
mate choice, 1554
Metarhizium anisopliae, 1554
reproductive maturation, 1554
soil moisture,
egg, 157
egg pod, 157
hatching, 157
oviposition, 157
Solanum viarum,
Florida hammocks, 1437
insect herbivory, 1437
invasive weed, 1437
pastures, 1437
Solenopsis invicta,
Decapitating fly, 73
diffusion model, 73
dispersal, 73
southern California,
ant community, 204
coastal sage scrub, 204
species diversity, 204
wildfire, 204

southern green stink bug,
brown stink bug, 1159
cotton, 1159
peanut, 1159
stink bug distribution, 1159
southern pine beetle,
cellular automata, 273
county pattern, 273
forest damage, 273
Spalangia cameroni,
Muscidifurax raptorellus, 88
parasitism, 88
Spalangia endius, 88
Stomoxys calcitrans, 88
spatial,
pheromone, 654
precision pest management, 654
Striacosta albicosta, 654
western bean cutworm, 654
spatial dispersion model,
distribution, 111
negative binomial distribution, 111
population ecology, 111
Sahlbergella singularis, 111
spatial distribution, 1253, 1511
Eurygaster integriceps, 1253
geostatistics, 1253
phenology, 1511
population, 1511
SADIE, 1253
Tortricidae, 1511
volatiles, 1511
spatial dynamics,
insect resistance management, 964
pest behavior, 964
refuge, 964
seed mixture, 964
spatial pattern,
ant mosaic, 999, 1067
ants, 999
pastures, 999
scramble competition, 999
self-organization, 1067
species composition, 231, 797
EMEND, 797
forest management, 797
invasive species, 231
population monitoring, 231
seasonality, 231
stink bug, 231
variable retention, 797
vertical stratification, 797
species diversity,
ant community, 204
coastal sage scrub, 204
southern California, 204
wildfire, 204
species interactions,
climate change, 824
community structure, 824
indirect interactions, 824
physiological ecology, 824
spiders,
consumption rate, 333
generalist insect predators, 333
killing rate, 333
natural enemies, 333
spined soldier bug,
developmental time, 948
predation, 948
yellow margined leaf beetle, 948
stable isotopes,
Bt corn, 1585
female longevity, 1585
head capsule width, 1585
male investment, 1585
starvation,
Coptotermes formosanus, 1232
moisture, 1232
mortality, 1232
sterile insect technique,
body size, 931
host, 931
Mediterranean fruit fly, 931
temperature, 931
sticky traps,
color response, 120
Elatobium abietinum, 120
flight activity, 120
Sitka spruce, 120
stink bug,
invasive species, 231
population monitoring, 231
seasonality, 231
species composition, 231
stink bug distribution,
brown stink bug, 1159
cotton, 1159
peanut, 1159
Southern green stink bug, 1159
Stomoxys calcitrans, 88, 184
larval development, 184
Muscidifurax raptorellus, 88
parasitism, 88
population dynamics, 184
Spalangia cameroni, 88
Spalangia endius, 88
stored grain,
developmental thresholds, 788
population growth, 788
stored products, 788
stored products,
developmental thresholds, 788
population growth, 788
stored grain, 788
streams,
freshwater, 289
microsporidia, 289
nematodes, 289
parasitism, 289
Striacosta albicosta,
pheromone, 654
precision pest management, 654
spatial, 654
western bean cutworm, 654
subarctic,
Acrididae, 541
bottom-up forces, 541
Orthoptera, 541
resource limitation, 541
succession,
disturbance, 1051
fire, 1051
insect, 1051
invertebrate, 1051
sugarcane borer,
alternate hosts, 1036
Diatraea saccharalis (F.), 1036
Eoreuma loftini (Dyar), 1036
Mexican rice borer, 1036
supercooling point,
cold hardiness, 1546
diapause, 1546
elm leaf beetle, 1546
Xanthogaleruca luteola, 1546
surveillance,
detection, 1276
food preference, 1276
foraging distance, 1276

model, 1276
susceptibility,
glassy-winged sharpshooter, 1192
grape, 1192
Pierce's disease, 1192
transmission efficiency, 1192
swallow-worts,
Chrysomelidae, 1427
host specificity, 1427
nontarget effects, 1427
prerelease impact, 1427
symbiont,
canola pest, 818
MLST, 818
multilocus sequence typing, 818
Wolbachia, 818
Synanthedon exitiosa,
Prunus angustifolia, 1465
Prunus persica, 1465
Prunus serotina, 1465
Synanthedon pictipes, 1465
Synanthedon pictipes,
Prunus angustifolia, 1465
Prunus persica, 1465
Prunus serotina, 1465
Synanthedon exitiosa, 1465
Synanthedon scitula,
apple, 173
geostatistics, 173
median polishing, 173
residuals, 173
syndrome 'basses richesses',
disease emergence, 1345
Hemiptera, 1345
vector, 1345

T

tangent equation,
Arrhenius plot, 462
enzyme inactivation, 462
intrinsic optimum temperature, 462
Sharpe-Schoolfield model, 462
Taylor's power law,
aggregation, 689
autocorrelation, 689
pest management, 689
Semivariogram, 689
Taylor's power law, 689
Telenomus podisi,
Euschistus servus, 303
Nezara viridula, 303
Trissolcus basalis, 303
temperature, 1, 931, 1577
body size, 1, 931
development, 1577
development time, 1
food, 1577
genotype-temperature interactions, 1
host, 931
Mediterranean fruit fly, 931
sterile insect technique, 931
Typhlodromus athenas, 1577
temperature requirement,
cotton aphid, 431
fecundity, 431
lower developmental threshold, 431
thermal constant, 431
temperature-size rule,
body size ratio, 737
foraging, 737
handling time, 737
Tephritidae,
Bactrocera dorsalis complex, 844
invasive fruit fly, 844
polyphagous, 844
quarantine pest, 844
tethered-flight,
Chrysoperla sinica, 449
migration, 449
ovarian development, 449
Tetropium,
Cerambycidae, 714
cinnamopterum, 714
contact pheromone, 714
fuscum, 714
thermal biology,
alfalfa leafcutting bee, 917
lipids, 917
Megachile rotundata, 917
pollinators, 917
thermal constant,
cotton aphid, 431
fecundity, 431
lower developmental threshold, 431
temperature requirement, 431
thermal threshold,
aphids, 1541
developmental rate, 939
field validation, 939
predator, 1541
pumpkin, 1541
rearing, 1541
variable temperature, 939
Xylotrechus arvicola, 939
thrips,
flight dispersal, 873
Mangifera indica, 873
yellow sticky trap, 873
Thrips tabaci,
Allium cepa, 194
population ecology, 194
Tospovirus, 194
tidal activity,
inundation, 1295
larval gas exchange, 1295
plastron, 1295
salt water, 1295
Tier-I toxicity assay,
dietary exposure assay, 1613
Ephestia egg, 1613
Orius insidiosus, 1613
plant-incorporated protectants (PIPs), 1613
tiered testing,
E-64, 496
environmental risk assessment, 496
nontarget effects, 496
potassium arsenate, 496
Tortricidae,
phenology, 1511
population, 1511
spatial distribution, 1511
volatiles, 1511
Tospovirus,
Allium cepa, 194
population ecology, 194
Thrips tabaci, 194
trait-meditated interactions,
agroforestry, 581
Azteca instabilis, 581
nonconsumptive effects, 581
Pseudacteon, 581
transgenic crops,
Bacillus thuringiensis, 484
feeding behavior, 484
refuge strategy, 484

simulation modeling, 484
transgenic GNA wheat,
EPG, 743
feeding behavior, 743
life table parameters, 743
wheat aphids, 743
transgenic rice,
Bt protein expression, 1323
control efficacy, 1323
ELISA, 1323
transmission efficiency,
glassy-winged sharpshooter, 1192
grape, 1192
Pierce's disease, 1192
susceptibility, 1192
trap crop,
natural refuge, 749
nonrice host plants, 749
organic crucifer production, 1471
pest management, 1471
resistance management, 749
tree condition,
Cerambycidae, 1200
development time, 1200
tree moisture stress, 1200
tree moisture stress,
Cerambycidae, 1200
development time, 1200
tree condition, 1200
trehalose,
glycogen, 1566
Gryllus firmus, 1566
triglyceride, 1566
wing dimorphism, 1566
Trialeurodes vaporariorum,
alien species, 140
Bemisia tabaci, 140
co-occurrence, 140
interspecific interaction, 140
Tribolium castaneum,
bioassay, 82
chemical attraction, 82
disturbance, 82
Trifolium pratense,
fatty acids, 399
Hylastinus obscurus, 399
red clover, 399
Scolytidae, 399
triglyceride,
glycogen, 1566
Gryllus firmus, 1566
trehalose, 1566
wing dimorphism, 1566
Trinidad,
Cuba, 989
Dominican Republic, 989
Guadeloupe, 989
Martinique, 989
Trissolcus basalis,
Euschistus servus, 303
Nezara viridula, 303
Telenomus podisi, 303
tritrophic interaction,
biological control, 260
ecosystem service, 260
tropical caterpillar,
insect-microbe interactions, 1111
midgut microbiota, 1111
Rothschildia lebeau, 1111
Área de Conservaci ón Guanacaste, 1111
Tsuga canadensis,
active hunters, 1378
habitat structure, 1378
predators, 1378
web builders, 1378
Tsuga spp.,
Adelges tsugae, 864
biological control, 864
host finding behavior, 864
predator-prey interactions, 864
Tunisia,
Bemisia tabaci, 809
secondary symbionts, 809
turfgrass,
golf courses, 589
introduction, 589
Ovavesicula popilliae, 589
Popillia japonica, 589
Typhlodromus athenas, 597, 1577
development, 597, 1577
food, 1577
life table, 597
Phytoseiidae, 597
pollen, 597
temperature, 1577

U

urban,
community ecology, 1397
habitat structure, 1397
landscaping, 1397
odorous house ant, 1397

V

Vaccinium corymbosum,
family diversity, 247
family richness, 247
herbivorous, 247
landscape, 247
Vaccinium corymbosum x *V. darrowi*,
Frankliniella bispinosa, 893
inverse distance weighting, 893
kriging, 893
semivariogram, 893
vacuum,
adult distribution, 661
codling moth, 661
diel behavior, 661
mating disruption, 661
variable retention,
EMEND, 797
forest management, 797
species composition, 797
vertical stratification, 797
variable temperature,
developmental rate, 939
field validation, 939
thermal threshold, 939
Xylotrechus arvicola, 939
vector,
disease emergence, 1345
Hemiptera, 1345
syndrome 'basses richesses', 1345
vertical stratification,
EMEND, 797
forest management, 797
species composition, 797
variable retention, 797
vineyards,
multiplex PCR, 1595
pest identification, 1595
Planococcus, 1595
Pseudococcus, 1595

virus sink,
Aphis gossypii, 391
barrier crop, 391
Myzus persicae, 391
nonpersistent virus, 391
visual,
host-finding behavior, 1494
huanglongbin, 1494
olfactory, 1494
plant–herbivore interactions, 1494
visual cues,
host finding, 855
host selection, 855
insect, 855
plant, 855
visual inspection,
binomial sampling, 381
classification sampling, 381
flower taps, 381
operating characteristic, 381
Vitacea polistiformis,
behavior, 880
plant-insect interactions, 880
roots, 880
volatiles, 126, 1511
cocoon, 126
frass, 126
Monodontomerus aeneus, 126
Osmia cornuta, 126
phenology, 1511
population, 1511
spatial distribution, 1511
Tortricidae, 1511
voltinism,
climate change, 505
grape berry moth, 505
greenhouse gas emissions, 505
phenology, 505

W

waste management,
Bacillus subtilis, 30
chicken manure, 30
Hermetia illucens, 30
web builders,
active hunters, 1378
habitat structure, 1378
predators, 1378
Tsuga canadensis, 1378
weed biological control,
Brazilian pepper, 605
exocrine gland, 605
host specificity testing, 605
western bean cutworm,
pheromone, 654
precision pest management, 654
spatial, 654
Striacosta albicosta, 654
western cherry fruit fly,
head width, 1353
Prunus avium, 1353
Prunus emarginata, 1353
wing length, 1353
western tarnished plant bug, 367, 455
cotton, 367
diapause, 455
feeding behavior, 367
Gossypium, 367
Lygus hesperus, 455
Lygus, 367
rearing, 455
western wheat aphid,
grass aphid, 151
greenbug, 151
pea aphid, 151
Russian wheat aphid, 151
wheat,
alternative host, 1303
gene-for-gene interaction, 1303
landscape genetics, 1303
microsatellites, 1303
wheat aphids,
EPG, 743
feeding behavior, 743
life table parameters, 743
transgenic GNA wheat, 743
wildfire,
ant community, 204
coastal sage scrub, 204
southern California, 204
species diversity, 204
wing dimorphism,
glycogen, 1566
Gryllus firmus, 1566
trehalose, 1566
triglyceride, 1566
wing length,
head width, 1353
Prunus avium, 1353
Prunus emarginata, 1353
Western cherry fruit fly, 1353
Winkler extractor,
Arachnida, 1523
Coleoptera, 1523
Formicidae, 1523
Micro-Hymenoptera, 1523
Wolbachia, 818, 1102
Burkholderia, 1102
canola pest, 818
Coreidae, 1102
crypts, 1102
gut symbiont, 1102
MLST, 818
multilocus sequence typing, 818
symbiont, 818
wood-borers,
Agrilus anxius, 648
biogeography, 648
coevolution, 648
host plant resistance, 648
woody hedgerow,
agro-ecosystems, 560
moth diversity, 560
organic farming, 560
plant diversity, 560

X

Xanthogaleruca luteola,
cold hardiness, 1546
diapause, 1546
elm leaf beetle, 1546
supercooling point, 1546
Xylella fastidiosa,
Aphrophoridae, 1131
Cicadellidae, 1131
Clastopteridae, 1131
Membracidae, 1131
Xylotrechus arvicola,
developmental rate, 939
field validation, 939
thermal threshold, 939
variable temperature, 939

Y

yellow margined leaf beetle,
developmental time, 948
predation, 948
spined soldier bug, 948
yellow mealworm,
behavior, 1285
dietary self-selection, 1285
insect rearing, 1285
nutrition, 1285
yellow sticky trap,
flight dispersal, 873
Mangifera indica, 873
thrips, 873

Z

(Z)-3-hexenyl acetate,
β-farnesene, 420
farnesol, 420
Monitoring, 420
pear ester, 420